




Draft Whole-Genome Sequence of a Uropathogenic *Escherichia coli* Strain Carrying the *eae* Gene

Tiago Barcelos Valiatti,^a Fernanda Fernandes Santos,^a Ana Carolina de Mello Santos,^a Rosa Maria Silva,^a Eneas Carvalho,^b  Tânia Aparecida Tardelli Gomes^a

^aDepartamento de Microbiologia, Imunologia e Parasitologia, Escola Paulista de Medicina, Universidade Federal de São Paulo, São Paulo, Brazil

^bLaboratório de Bacteriologia, Instituto Butantan, São Paulo, Brazil

ABSTRACT Uropathogenic *Escherichia coli* (UPEC) strains are responsible for most cases of urinary tract infections worldwide. We present the draft whole-genome sequence of the UPEC 252 strain, which carries the *eae* gene that encodes the intimin adhesin. Intimin promotes intimate adherence of enteropathogenic *E. coli* and enterohemorrhagic *E. coli* to intestinal cells.

Urinary tract infection (UTI) is considered one of the most frequent infections affecting the community and hospitalized individuals worldwide (1). The main causative agent of UTI is uropathogenic *Escherichia coli* (UPEC), which represents about 75 to 90% of cases (1–3).

This announcement aims to describe a draft whole-genome sequence of UPEC 252, isolated in 1999 from a 1-month-old female patient with a UTI who was admitted to Hospital São Paulo, a public hospital in São Paulo, Brazil. The UPEC 252 strain was originally characterized as a carrier of the *eae* gene (4), which encodes intimin, the main adhesin involved in the ability of enteropathogenic *E. coli* (EPEC) and enterohemorrhagic *E. coli* (EHEC) to promote attaching and effacing (AE) lesions in epithelial cells (5).

UPEC 252 was stored in lysogeny broth (LB) plus 15% glycerol and kept at –80°C. One isolated colony obtained on MacConkey agar was cultured overnight in LB at 37°C, and DNA extraction was performed using the Wizard genomic DNA purification kit (Promega USA), following the manufacturer's recommendations. Genomic DNA was fragmented by tagmentation using the Nextera XT DNA library preparation kit, according to the manufacturer's instructions, yielding fragments with an average size of 1 kb. A genomic library was prepared using the HiSeq rapid SBS v2 library kit (Illumina USA), which was subsequently sequenced on the Illumina HiSeq 1500 platform using the 2 × 250-bp protocol. The total number of reads sequenced was 5,267,541, which was reduced to 4,876,002 reads (92.5%) after the removal of low-quality bases (regions with a mean quality score below 25 were discarded), small reads (shorter than 40 bases), and Illumina adaptors using the software Trimmomatic (6). The data that originated from the Illumina readings were assembled using SPAdes version 3.12.0 (7), annotation was performed using Prokka version 1.13.3 (8), and the genome assembly metric was calculated using QUAST. All software programs were used with the default settings.

The genome of UPEC 252 presented a total size of 4,802,947 bp (GC content, 50.46%) distributed in 151 contigs, the largest being 322,751 bp. Further, the N_{50} and N_{75} values were 187,460 and 87,691 bp, respectively, while the L_{50} and L_{75} values were 10 and 20 contigs, respectively. The annotation showed that UPEC 252 has 1 transfer-messenger RNA (tmRNA) gene, 83 tRNA genes, 8 rRNA genes, and 4,520 coding sequences (CDS).

The contigs were submitted to the VirulenceFinder software (version 2.0), Center for Genomic Epidemiology (9), to identify the genes belonging to the pathogenicity island of approximately 35 kb called the locus of enterocyte effacement (LEE). The LEE

Citation Valiatti TB, Santos FF, Santos ACDM, Silva RM, Carvalho E, Gomes TAT. 2019. Draft whole-genome sequence of a uropathogenic *Escherichia coli* strain carrying the *eae* gene. *Microbiol Resour Announc* 8:e00980-19. <https://doi.org/10.1128/MRA.00980-19>.

Editor Irene L. G. Newton, Indiana University, Bloomington

Copyright © 2019 Valiatti et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Fernanda Fernandes Santos, ff.santos@unifesp.br, or Tânia Aparecida Tardelli Gomes, tatg.amaral@unifesp.br.

Received 12 August 2019

Accepted 1 October 2019

Published 24 October 2019

includes the *eae* gene, which is required for AE lesion formation, suggesting that UPEC 252 has hybrid potential, containing virulence genes associated with the establishment of diarrhea and urinary tract infections.

We expect that knowledge of the UPEC 252 genome will contribute to the support of future studies on the true virulence potential of hybrid pathogenic *E. coli* strains.

Data availability. The reads used for assembly of the UPEC 252 genome were deposited in the Sequence Read Archive (SRA) at the NCBI under the accession number [SRR9317828](https://www.ncbi.nlm.nih.gov/sra/SRR9317828), and the whole-genome shotgun sequences were deposited in the GenBank database under the accession number [VFST00000000](https://www.ncbi.nlm.nih.gov/genbank/VFST00000000). The versions described in this paper are the first versions.

ACKNOWLEDGMENT

This study was conducted with a scholarship funded by grant 2017/21947-7 to T.B.V. and grant 2018/17353-7 to T.A.T.G. from the São Paulo Research Foundation (FAPESP).

REFERENCES

- Hooton T. 2012. Uncomplicated urinary tract infection. *N Engl J Med* 366:1028–1037. <https://doi.org/10.1056/NEJMcp1104429>.
- Korbel L, Howell M, Spencer JD. 2017. The clinical diagnosis and management of urinary tract infections in children and adolescents. *Paediatr Int Child Health* 37:273–279. <https://doi.org/10.1080/20469047.2017.1382046>.
- Ronald A. 2003. The etiology of urinary tract infection: traditional and emerging pathogens. *Dis Mon* 49:71–82. <https://doi.org/10.1067/mda.2003.8>.
- Abe CM, Salvador FA, Falsetti IN, Vieira MA, Blanco J, Blanco JE, Blanco M, Machado AM, Elias WP, Hernandez RT, Gomes TA. 2008. Uropathogenic *Escherichia coli* (UPEC) strains may carry virulence properties of diarrhoeagenic *E. coli*. *FEMS Immunol Med Microbiol* 52:397–406. <https://doi.org/10.1111/j.1574-695X.2008.00388.x>.
- Kaper JB, Nataro JP, Mobley HL. 2004. Pathogenic *Escherichia coli*. *Nat Rev Microbiol* 2:123–140. <https://doi.org/10.1038/nrmicro818>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- Joensen KG, Scheutz F, Lund O, Hasman H, Kaas RS, Nielsen EM, Aarestrup FM. 2014. Real-time whole-genome sequencing for routine typing, surveillance, and outbreak detection of verotoxigenic *Escherichia coli*. *J Clin Microbiol* 52:1501–1510. <https://doi.org/10.1128/JCM.03617-13>.