

DMEM high glucose and polystyrene surface

Biofilm formation was measured as absorbance at 595 nm, with 1% α -D-man

Repetitions	EAEC 042					UPEC-46					<i>E. coli</i> DH5 α				
	3 h	6 h	9 h	12 h	24 h	3 h	6 h	9 h	12 h	24 h	3 h	6 h	9 h	12 h	24 h
A(1)	0	1.237	2.001	2.342	2.534	0	0	0	0	0.178	0	0	0.089	0.023	0.158
A(2)	0	1.897	2.124	2.654	2.678	0	0	0	0.058	0.214	0	0	0.012	0	0.14
A(3)	0.102	1.987	2.076	2.456	2.875	0	0	0	0.079	0.198	0	0	0.067	0	0.121
B(1)	0.045	2.001	2.231	2.756	3	0	0	0	0.124	0.153	0	0	0	0.012	0.238
B(2)	0.069	1.456	2.087	2.421	2.432	0	0	0	0.102	0.149	0	0	0	0.097	0.181
B(3)	0	1.678	1.945	2.354	2.298	0	0	0	0.1	0.123	0	0	0	0.056	0.158
C(1)	0.123	1.862	1.921	2.012	2.765	0	0	0	0.078	0.243	0	0	0.101	0.121	0.128
C(2)	0.111	2.1	2.1	2.134	2.543	0	0	0	0.046	0.222	0	0	0.124	0.049	0.107
C(3)	0.012	1.965	2.021	2.132	2.431	0	0	0	0.05	0.218	0	0	0.098	0.109	0.133

Note: The assays were performed in triplicate and repeated three times