

DMEM high glucose and polystyrene surface
Biofilm formation was measured as absorbance at 595 nm, without α -D-man

Repetitions	EAEC 042					UPEC-46					<i>E. coli</i> DH5 α				
	3 h	6 h	9 h	12 h	24 h	3 h	6 h	9 h	12 h	24 h	3 h	6 h	9 h	12 h	24 h
A(1)	0.275	2.589	2.594	2.434	2.265	0.257	0.386	0.628	0.543	0.288	0	0	0.006	0.098	0.106
A(2)	0.312	2.413	2.439	2.453	2.303	0.217	0.518	0.693	0.601	0.287	0	0	0.01	0.1	0.201
A(3)	0.389	2.531	2.969	2.534	2.464	0.129	0.361	0.534	0.598	0.323	0	0	0.1	0.102	0.1567
B(1)	0.432	2.214	2.654	2.456	2.543	0.213	0.301	0.7	0.567	0.432	0	0	0.101	0.123	0.165
B(2)	0.435	2.087	2.876	2.654	2.21	0.278	0.301	0.623	0.678	0.398	0	0	0.088	0.121	0.165
B(3)	0.234	2.189	2.545	2.654	2.345	0.201	0.231	0.543	0.421	0.423	0	0	0.056	0.111	0.189
C(1)	0.567	2.67	2.67	2.564	2.543	0.126	0.412	0.674	0.398	0.436	0	0	0	0.109	0.1654
C(2)	0.512	2.312	2.789	2.345	2.523	0.163	0.345	0.611	0.301	0.245	0.003	0	0	0.132	0.168
C(3)	0.486	2.123	2.754	2.479	2.234	0.189	0.399	0.622	0.467	0.345	0.009	0	0	0.123	0.175

Note: The assays were performed in triplicate and repeated three times