

Human urine and polystyrene surface
Biofilm formation was measured as absorbance at 595 nm, with 1% α -D-man

Repetitions	EAEC 042					UPEC-46					<i>E. coli</i> DH5 α				
	3 h	6 h	9 h	12 h	24 h	3 h	6 h	9 h	12 h	24 h	3 h	6 h	9 h	12 h	24 h
A(1)	0.06	0.468	0.307	0.35	0.54	0.005	0.005	0.03	0.008	0.017	0	0.002	0.008	0.015	0.027
A(2)	0.05	0.395	0.262	0.381	0.675	0.002	0.003	0.01	0.026	0.043	0	0	0	0.011	0.04
A(3)	0.022	0.391	0.25	0.294	0.547	0	0	0.042	0.032	0.014	0	0	0	0.018	0.028
B(1)	0.078	0.447	0.212	0.439	0.413	0	0	0.041	0.032	0.045	0.002	0	0.019	0	0.044
B(2)	0.069	0.385	0.416	0.395	0.674	0	0.002	0.038	0.006	0.038	0	0	0.014	0	0.025
B(3)	0.058	0.433	0.378	0.449	0.45	0.025	0.017	0.028	0.048	0.017	0.006	0	0.027	0.001	0
C(1)	0.041	0.372	0.324	0.461	0.54	0.023	0	0.024	0.015	0.007	0.014	0	0.01	0	0.038
C(2)	0.027	0.305	0.436	0.558	0.571	0.029	0.009	0.023	0.008	0.042	0	0.004	0.017	0	0.005
C(3)	0.108	0.369	0.325	0.501	0.496	0.029	0.015	0.018	0.02	0.032	0.006	0.006	0.001	0	0.061

Note: The assays were performed in triplicate and repeated three times