

**Quantitative adherence assays of UPEC-46 strain in absence of D-mannose expressed in CFU/mL**

<b>Repetitions</b>	<b>HeLa</b>	<b>HT-29</b>	<b>5637</b>
A(1)	6000000	97000000	100000000
A(2)	8000000	80000000	100000000
B(1)	15000000	63000000	10000000
B(2)	10000000	100000000	10000000
C(1)	12000000	130000000	20000000
C(2)	10000000	40000000	33000000

Note: 3 h of incubation. The assays were performed in duplicate and repeated three times

**Quantitative adherence assays of UPEC-46 strain in absence of D-mannose expressed in Log<sub>10</sub> CFU/mL**

<b>Repetitions</b>	<b>HeLa</b>	<b>HT-29</b>	<b>5637</b>
A(1)	6.778	7.987	8
A(2)	6.903	7.903	8
B(1)	7.176	7.799	7
B(2)	7	8	7
C(1)	7.079	8.114	7.301
C(2)	7	7.602	7.519

Note: 3 h of incubation. The assays were performed in duplicate and repeated three times

**Quantitative adherence assays of UPEC-46 strain in persence of 1% D-mannose expressed in CFU/mL**

<b>Repetitions</b>	<b>HeLa</b>	<b>HT-29</b>	<b>5637</b>
A(1)	6000000	13000000	4000000
A(2)	4700000	17000000	4300000
B(1)	4000000	9000000	5000000
B(2)	7700000	9700000	5300000
C(1)	1700000	10000000	1000000
C(2)	3000000	5000000	10000000

Note: 3 h of incubation. The assays were performed in duplicate and repeated three times

**Quantitative adherence assays of UPEC-46 strain in persence of 1% D-mannose expressed in Log<sub>10</sub> CFU/mL**

<b>Repetitions</b>	<b>HeLa</b>	<b>HT-29</b>	<b>5637</b>
A(1)	6.778	7.114	6.602
A(2)	6.672	7.23	6.633
B(1)	6.602	6.954	6.699
B(2)	6.886	6.987	6.724
C(1)	6.23	7	6
C(2)	6.477	6.699	7

Note: 3 h of incubation. The assays were performed in duplicate and repeated three times