




Taxonomic revision and additional comments of some bats (Mammalia, Chiroptera) reported from Bolivia, with an updated checklist based on voucher material with verified identities

JOSÉ L. POMA-UREY^{1*}, LUIS H. ACOSTA S.¹, KATHIA RIVERO¹, MARISOL HIDALGO-COSSIO², ERIKA HINGST-ZAHER³, JULIANA GUALDA-BARROS⁴, BEATRIZ D. DA NATIVIDADE⁵, KATHRIN BARBOZA-MARQUEZ¹, HÉCTOR E. RAMÍREZ-CHAVES⁶, JORGE SALAZAR-BRAVO⁷, JOSÉ OCHOA G.⁸


¹ Área de Zoología, Museo de Historia Natural Noel Kempff Mercado de la Universidad Autónoma Gabriel René Moreno, Santa Cruz, Bolivia • JLP: jose_luispoma@hotmail.com • LHAS: ljubatus096@gmail.com • KR: krivero@museoelkempff.org • KBM: batikathrincita@gmail.com


² Colección Boliviana de Fauna, Museo Nacional de Historia Natural de Bolivia; Área de Mastozoología, Museo de Historia Natural Alcide d'Orbigny, Cochabamba, Bolivia • MHC: hidalgocossiomarisol@gmail.com

³ Museu Biológico, Instituto Butantan, São Paulo, SP, Brazil • EH: erika.zaher@butantan.gov.br  <https://orcid.org/0000-0001-9547-6666>

⁴ Laboratório de Mastozoologia, Museu de Zoologia da Universidade São Paulo, São Paulo, SP, Brazil • JGB: jgbarros@usp.br  <https://orcid.org/0000-0003-2443-0788>

⁵ Departamento de Sistemática e Ecologia, Laboratório de Mamíferos, Universidade Federal da Paraíba, João Pessoa, PB, Brazil • BDN: bdnatividade@gmail.com

⁶ Departamento de Ciencias Biológicas, Centro de Museos, Museo de Historia Natural, and Natural History Laboratory Integrative Zoological Biodiversity Discovery, Universidad de Caldas, Manizales, Caldas, Colombia • HERC: hector.ramirez@ucaldas.edu.co  <https://orcid.org/0000-0002-2454-9482>

⁷ Department of Biological Sciences, Texas Tech University, Lubbock, TX, USA • Instituto Nacional de Biodiversidad, Quito, Ecuador • JSB: j.salazarbravo@ttu.edu  <https://orcid.org/0000-0001-5905-2532>

⁸ Centro de Estudios en Zoología Aplicada, Universidad de Carabobo, Valencia, Venezuela • JOG: jochoabat@gmail.com

* Corresponding author

Abstract. We review the taxonomic identities of museum specimens of three bat species previously reported from Bolivia. We comment on some erroneously reported taxa, or taxa either not represented by voucher materials or based on insufficient data (including acoustic detections) to verify their taxonomic affinities. As result of this review, the list of bat species known to occur in Bolivia is updated to eight families and 133 species, unlike the nine families and 146 species of previous lists. Some recommendations for future research and a brief historical revision of bat inventories in the country are included.

Keywords. Biological collections, geographic distribution, invalid records, South America

Academic editor: Annia Rodríguez-San Pedro

Received 16 February 2023, accepted 14 June 2023, published 21 June 2023

Poma-Urey JL, Acosta S. LH, Rivero K, Hidalgo-Cossio M, Hingst-Zaher E, Gualda-Barros J, da Natividade BD, Barboza-Marquez K, Ramirez-Chaves HE, Salazar-Bravo J, Ochoa G. J (2023) Taxonomic revision and additional comments of some bats (Mammalia, Chiroptera) reported from Bolivia, with an updated checklist based on voucher material with verified identities. *Check List* 19 (3): 409–427. <https://doi.org/10.15560/19.3.409>

Introduction

Knowledge of the diversity of Bolivian mammals is the result of numerous national and international research initiatives, mostly linked to faunal surveys and the management of systematic collections (Anderson 1997; Tarifa 2014). Between the 1960s and 2000, five expeditions were conducted, which made substantial contributions to the inventory of bats in the country. The first expedition, in 1963, was part of a project aimed at the diagnosis of hemorrhagic fever and the identification of its reservoirs in the Beni Department, mainly carried out by the Panama-based and US-sponsored Middle America Research Unit. Between 1963 and 1966, the American Museum of Natural History (New York, USA) organized the second expedition, whose objective was to recognize ectoparasites associated with wild mammals. A third expedition, under the responsibility of Louisiana State University Museum of Zoology (1979–1981), surveyed bats mainly in La Paz Department. In 1986, the fourth expedition was executed by the Zoológico Municipal of Santa Cruz, in the Santa Cruz Department, under the leadership of Professor Noel Kempff Mercado and with the participation of researchers from Doñana Biological Station (Seville, Spain) (Cabot-Nieves 1996; Anderson 1997; Tarifa 2014). Finally, between 1984 and 1996, an important joint research was conducted by the American Museum of Natural History, the Museum of Southwestern Biology (University of New Mexico, New Mexico), the Colección Boliviana de Fauna (La Paz), and the Museo “Noel Kempff Mercado” (Santa Cruz), which was fundamental in training Bolivian biologists, in addition to collecting an important number of museum specimens (not only bats) in a geographic area defined by hundreds of localities across the country (Anderson 1997; Tarifa 2014). These expeditions provided a taxonomic list that includes around 40% of the bat species known from Bolivia, and their results represents an important advancement for the knowledge of this group of mammals in several regions of the country.

The first checklists of bats collected in Bolivia were published by Anderson et al. (1982) and Anderson (1997), who reported the presence of 79 and 105 species grouped into five and seven families, respectively. Subsequently, Salazar-Bravo et al. (2003) updated the inventory with 109 species belonging to eight families, Aguirre (2007) compiled 122 species representing eight families, and, three years later, Aguirre et al. (2010) reported 132 species grouped into nine families, substantially increasing the number of taxa recorded in previous years. Recently, Aguirre et al. (2019) and Díaz et al. (2021) provided two additional lists of the bats of the country, in which 138 and 146 species were included, respectively.

The most recent contributions to the knowledge of the Bolivian bat fauna represent the results of several initiatives conducted by local researchers interested in taxonomic and ecological studies. Their investigations have

led to the discovery of undescribed species (e.g., Siles et al. 2013; Acosta et al. 2021) and the inclusion of additional species for the country (Calderón-Acevedo and Muchhala 2020; Poma-Urey et al. 2021). On the other hand, these initiatives made it possible to exclude species previously listed from Bolivia due to misidentifications (e.g., *Anoura fistulata* Muchhala, Mena and Albuja, 2005; Calderón-Acevedo and Muchhala 2018) or whose voucher material is not available for revision (e.g., *Micronycteris schmidtorum* Sanborn, 1935) (Aguirre et al. 2019).

The main principle and criteria supporting a taxonomic inventory of bat species of Bolivia is based on the golden rule that at least one voucher specimen of each taxon must be deposited in an open-access systematic collection for a species to be considered present in the country. In some cases, the unequivocal vocal signatures of a group of insectivorous bats (e.g., *Molossops* spp. and *Promops* spp.) allow to use their acoustic records as voucher material. However, in many other cases, there can be times when it is difficult to use echolocation calls to precisely identify species; examples of these situations are when bats are flying in clutter conditions or during social interactions, variations of the position or distance of bats with respect to the detector, and the absence of verified signals in a taxonomic context (Miller and Corben pers. comm.). Vouchers are the evidence for the occurrence of a species at a given time or place and constitute the baseline data on a species' distribution, habitat, morphological characteristics, echolocation patterns, and other aspects of the natural history (Ruedas et al. 2000; López-Vidal and Elizalde-Arellano 2006; Jung et al. 2007, 2014). In addition, because studies on the systematics of Neotropical bats are ongoing (e.g., Basantes et al. 2020; Esquivel et al. 2022), vouchered museum specimens are fundamental references of the taxonomic status and nomenclature changes of species.

In this study, we update the previous inventories of bats in Bolivia (Aguirre et al. 2019; Díaz et al. 2021), based on the existence of voucher specimens deposited in open-access systematic collections and the revision of their taxonomic identity. We exclude from our list species without available or verifiable voucher material, including those whose records based only on acoustic methods without fundamental evidence that allows for confirming their identities (e.g., reference images of typical search phase calls or the descriptions of basic parameters). In addition, we provide ancillary information for a handful of poorly documented species, in terms of their morphometric and qualitative characters (Emmons et al. 2006; Aguirre et al. 2010). Finally, we recognize by name a group of individuals and institutions whose contributions, either through of field and/or museum surveys, have generated an important part of the knowledge about the bats of Bolivia.

Methods

We reviewed systematic collections and literature for information validating the existence of voucher

material for each species included in previous list of bats recorded for Bolivia. The following institutions provided the data on museum specimens reported in this study: Centro de Biodiversidad y Genética (CBG), Cochabamba, Bolivia; Colección Boliviana de Fauna (CBF), La Paz, Bolivia; Museo Noel Kempff Mercado (MNKM), Santa Cruz, Bolivia; Museu de Zoologia da Universidade de São Paulo (MZUSP), São Paulo, Brazil and Universidade Federal da Paraíba (UFPB), João Pessoa, Brazil. Other valid literature records are found in the following systematic collections: American Museum of Natural History (AMNH), New York, USA; The Natural History Museum (BMNH), London, England; Carnegie Museum of Natural History (CM), Pittsburgh, USA; Colección de Mamíferos Lillo (CML), Tucuman, Argentina; Estación Biológica Doñana (EBD), Sevilla, Spain; Field Museum of Natural History (FMNH), Chicago, USA; Louisiana State University, Museum of Zoology (LSUMZ), Baton Rouge, USA; Museo de Historia Natural Alcide d'Orbigny (MHNC-M), Cochabamba, Bolivia; Michigan State University (MSU), East Lansing, USA; Museum of Texas Tech University (TTU-M), Lubbock, USA; National Museum of Natural History (USNM), Washington DC, USA; Universidade Estadual Paulista, São José do Rio Preto (SJRP), Brazil; University of Michigan, Museum Zoology (UMMZ), Ann Arbor, USA.

The data compiled from voucher specimens included collection name, catalogue number, locality (coordinates and elevation, when available), and the morphological features that allow systematic identification. When some of these items were not available, we add-

ed here information, including some somatic and dental skull measurements obtained following the criteria proposed by Barquez et al. (1999). On the base of this protocol, we generate an emended checklist of the bats from Bolivia, with the names of collectors and the dates in which each species was reported for the first time in the country. Importantly, this revised checklist is updated to 1 March 2023.

Results

We found a series of voucher specimens that have been misidentified in previous lists of bats from Bolivia and whose characters were re-evaluated, and their taxonomic affiliations are updated here. They are: Phyllostomidae: *Glyphonycteris behnii* (Peters, 1865), now assigned to *Micronycteris hirsuta* (Peters, 1869); Furipteridae: *Furipterus horrens* (Cuvier, 1828), now assigned to *Peropteryx macrotis* (Wagner, 1843), family Emballonuridae; and Vespertilionidae: *Histiotus macrotus* (Poeppig, 1835), now assigned to *H. diaphanopterus* Feijó, da Rocha & Althoff, 2015. Consequently, *G. behnii*, *F. horrens*, and *H. macrotus*, as well as the family Furipteridae, were eliminated from the list of bats known in Bolivia (Table 1).

We also examined the morphological characters of additional voucher materials of two species of Emballonuridae—*Cyttarops alecto* Tomas, 1913 and *Saccopteryx canescens* Thomas, 1901—known in Bolivia by a scarce number of records. These specimens represent significant distributional extensions and provide complementary data on the ecological patterns and taxonomy of both taxa.

Table 1. Wrong or unconfirmed original records, in addition to synonymized species previously reported from Bolivia. Acoustic records included in the list were not supported by fundamental data to confirm their identities (images of typical search phase calls or the description of basic parameters).

Taxa	References	Remarks
Emballonuridae (4)		
<i>Centronycteris maximiliani</i>	Venzal et al. 2013	Repository data or diagnostic characters not provided
<i>Cormura brevirostris</i>	Vargas-Espinoza 2007	Recorded with acoustic methods only
<i>Diclidurus albus</i>	Emmons et al. 2006	Recorded with acoustic methods only
<i>Peropteryx kappleri</i>	Emmons et al. 2006	Recorded with acoustic methods only
Phyllostomidae (5)		
<i>Anoura aequatoris</i>	Calderón-Acevedo et al. 2021, 2022	Junior synonym of <i>A. caudifer</i>
<i>Anoura fistulata</i>	Aguirre et al. 2019	No voucher specimen in systematic collection
<i>Anoura peruana</i>	Calderón-Acevedo et al. 2021, 2022	Junior synonym of <i>A. geoffroyi</i>
<i>Glyphonycteris behnii</i>	Terán 2010; this study	Specimen reidentified as <i>Micronycteris hirsuta</i>
<i>Micronycteris microtis</i>	Porter et al. 2007; Morales-Martínez et al. 2021	Junior synonym of <i>M. megalotis</i>
Furipteridae (1)		
<i>Furipterus horrens</i>	Aguirre et al. 2010; this study	Specimen reidentified as <i>Peropteryx macrotis</i>
Molossidae (2)		
<i>Molossus currentium</i>	Moya et al. 2007	Recorded with acoustic methods only
<i>Molossus pretiosus</i>	Aguirre et al. 2010; this study	No voucher specimen in systematic collection
Vespertilionidae (1)		
<i>Histiotus macrotus</i>	Acosta and Vanegas 2006; this study	Specimen reidentified as <i>H. diaphanopterus</i>



Figure 1. External details of *Micronycteris hirsuta*, CBF 10666. **A** dorsal fur. **B, C.** Tuft of hair and band of skin connecting the ears.

As a complement to the results described above, we found that six species previously included in the inventories of bat species known from Bolivia are not represented by verifiable vouchers in any systematic collection and some of them have been recorded only by acoustic detection without supporting data that allows for their identities to be confirmed (i.e., reference images of typical search phase calls or descriptions of fundamental parameters; Table 1). This group of taxa includes *Centronycteris maximiliani* (Fischer, 1829) (Venzal et al. 2013), *Cormura brevirostris* (Wagner, 1843) (Vargas-Espinoza 2007), *Diclidurus albus* Wied-Neuwied, 1820, *Peropteryx kappleri* Peters, 1867 (Emmons et al. 2006), *Molossus currentium* Thomas, 1901 (Moya et al. 2007), and *M. pretiosus* Miller, 1902 (Aguirre et al. 2010). In addition, some specimens suspected to represent *P. kappleri* (Aguirre and Urioste 1994) are *P. leucoptera* Peters, 1867 (Poma-Urey et al. 2021), and three taxa have been incorporated into synonymies: *Anoura aequatoris* (Lönnerberg, 1921), *A. peruana* (Tschudi, 1844), and *Micronycteris microtis* Miller, 1898 (Porter et al. 2007; Morales-Martínez et al. 2021; Calderón-Acevedo et al. 2021, 2022). Finally, *Anoura fistulata*, considered by Aguirre et al. (2019) as member of the bat fauna of Bolivia, is not represented by a voucher specimen to verify the presence of this taxon in

the country and one specimen previously identified as *A. fistulata* (FMNH 106088; Mantilla-Meluk et al. 2014) is *A. caudifer* (É. Geoffroy, 1818) (Calderón-Acevedo and Muchhala 2018).

Order Chiroptera
Family Phyllostomidae

***Micronycteris hirsuta* (Peters, 1869)**

Material examined. BOLIVIA – La Paz • Ixiamas; 13°59'02"S, 068°52'25"W; 1492 m alt.; 25.III.2003; Marcos Terán leg.; 1 ♂ adult, CBF 10666 (Figs. 1A–C, 2)

Identification. The specimen matches the qualitative descriptions of *Micronycteris* Gray, 1866 (Fig. 1), but it was erroneously identified as *Glyphonycteris behnii* by Terán (2010). Bats of the genus *Micronycteris* are characterized by having bicolored dorsal hair with a white base and brown tip, distally rounded ears joined by a band of skin across the head (Fig. 1B, C), and a chin with a pair of V-shaped dermal pads without a central papilla. In contrast, members of the genus *Glyphonycteris* Thomas, 1896 are mainly characterized by presenting unicolor or tricolor dorsal hair, but not bicolored, in addition to lacking a band of skin on the head that connects the ears (Simmons and Voss 1998). Some measurements of the specimen examined are as follows:




Figure 2. Geographic distribution in Bolivia of the revised specimens of the following species: 1 = *Cyttarops alecto*, CBG 401; 2 = *Miconycteris hirsuta*, CBF 10666; 3 = *Saccopteryx canescens*, MNKM 1910; 4 = *Histiotus diaphanopterus*, MNKM 3713; 5 = *Peropteryx macrotis*, MZUSP 7733.

total length = 67 mm, tail length = 13 mm, foot length = 13 mm, ear length = 24 mm, calcar length = 15 mm, forearm length = 47.36 mm, and weight = 15.5 g. The combination of qualitative characters and measurements of this specimen match with the diagnosis of *M. hirsuta*, which is the largest species among its congeners.

Remarks. The specimen CBF 10666 was erroneously considered as the only record of *G. behnii* from Bolivia by Terán (2010), who did not indicate the systematic collection where the voucher specimen was deposited nor any diagnostic characters by which it was assigned to *G. behnii*. However, this specimen was recently incorporated into the Bolivian Fauna Collection (CBF

10666), where a detailed morphological examination allowed us to identify it as *M. hirsuta*. This is only the second record of this bat species for Bolivia (Fig. 2), after the report of Azurduy and Emmons (2005). Considering that Terán's (2010) record of *G. behnii* from Bolivia was misidentified, the species maintains its status as endemic to Brazil (Zortea et al. 2016; Solari et al. 2019; Burgin et al. 2020).

Family Emballonuridae

Peropteryx macrotis (Wagner, 1843)

Material examined. BOLIVIA – Santa Cruz • San José de Chiquitos; 17°51'S, 060°47'W; 311 m alt; V.1954; Carl Gans leg.; 1 ♀ adult, MZUSP 7733 (Figs. 2, 3A, B).




Figure 3. Dorsal and ventral views of external traits of some bat specimens. **A, B.** *Peropteryx macrotis* (MZUSP 7733). **C, D.** *Furipterus horrens* (UFPB 7994). Note free thumbs in *P. macrotis* vs. thumbs covered in *F. horrens*. The black arrow indicates the free tail on the uropatagium, while the red arrows indicate the phalanges of the third finger arranged first outward and then inward. **E, F.** View of the bicolored dorsal fur and the band joining the ears in *Histiotus diaphanopterus* (MNKM 3713), in addition to the highly developed lobe on the inner edge of ears (white arrow).

Family Furipteridae

Furipterus horrens (Cuvier, 1828)

Materials examined. BRAZIL – **Minas Gerais** • Rio Itacambirucu; 16°34'32"S, 041°24'17"W; 239 m alt.; 2.V.1989; Renato N. Feio leg.; 1 ♂ adult, UFPB 7994 (Fig. 3C, D) – **Bahia** • Complexo eólico Zeus; 10°29'13"S, 040°29'51" W; 822 m alt.; 29.III.2019; Mónica Pedrosa leg.; 1 ♂ adult, UFPB 10797 – **Paraná** • Colombo, Gruta do Bacaetava; 25°13'55"S, 049°12'27" W; 978 m alt.; 29.V.2015; Jennifer Barros leg.; 1 ♂ adult, UFPB 2185.

Identification. The specimen MZUSP 7733 was erroneously assigned to *Furipterus horrens* (Furipteridae) by Aguirre et al. (2010), even though its morphological characters correspond to the genus *Peropteryx* Peters, 1867 (Emballonuridae). In *F. horrens*, the forearm length is <38.5 mm, the tail is located within the uropatagium, and the thumbs are covered by the propatagium (Fig. 3C, D; Uieda et al. 1980; Novaes et al. 2012; Gardner 2008; Alfaro-Lara et al. 2018; Arroyo-Cabrales 2019). In contrast, MZUSP 7733, a female, has a forearm length of 42.76 mm, the phalanges of the third finger are arranged first outward and then inward, the tail is free over the uropatagium, the thumb is not covered by the propatagium, and the plagiopatagium is attached to the end of the tibia (Fig. 3A, B). Unfortunately, the skull is missing, and the forearm membrane is in poor condition, which prevents observation of the glandular sac; therefore, we considered other body parts when identifying this specimen. In considering the lack of ornamentation on the forearm and lines on the dorsal fur, we ruled out the genera *Rhynchonycteris* Peters, 1867 and *Saccopteryx* Illiger, 1811, respectively (although

S. gymnura has no lines but has a forearm length <36 mm). On the other hand, as the thumb is exposed and the fur is dark brown, the specimen does not belong to the genera *Cyttarops* Thomas, 1913 or *Diclidurus* Wied-Neuwied, 1820 (Hood and Gardner 2008). The specimen also does not represent the genus *Centronycteris* Gray, 1838, as the color of the latter ranges from orange to reddish and the plagiopatagium is attached to the metatarsals near the toes. Finally, it differs from *Cormura* Peters, 1867 in having the plagiopatagium not attached to the metatarsals at the base of the feet (Hood and Gardner 2008). Taking into account these characteristics, we infer that MZUSP 7733 belongs to the genus *Peropteryx* and, considering that it is of medium size without translucent wing membranes, that the species is *P. macrotis*.

Taxonomic remarks. The family Furipteridae was included as part of the bat fauna of Bolivia based on the single specimen examined here, erroneously identified as *F. horrens* (Aguirre et al. 2010) and currently deposited in the Museum of the University of São Paulo. Comparison of this specimen with three individuals of *F. horrens* from Brazil (UFPB 2185, 7994, 10797; Fig. 3C, D) confirms our revised identification; therefore, the family is excluded from the list of Bolivian bats.

Family Vespertilionidae

Histiotus diaphanopterus Feijó, da Rocha & Althoff, 2015

Materials examined. BOLIVIA – **Santa Cruz** • Vallegrande, Potrerillos; 18°33'S, 064°56'W; 2367 m alt.; 23.IV.1993; Israel Vargas leg.; 1 ♀ adult, MNKM 3713 (Figs. 2, 3E, F) • Florida, Pampagrande; 17°55'47"S, 064°09'




Figure 4. Lateral and dorsal views of the skulls of some bat species. **A.** *Cyttarops alecto* (CBG 401). **B.** *Saccopteryx canescens* (MNKM 1910). **C.** *Saccopteryx leptura* (MNKM 1908).

48°W; 1620 m alt.; I.1996; Arturo Muñoz leg.; 1 ♀ adult, MNKM 3714.

Identification. Acosta and Vanegas (2006) assigned to the female specimen MNKM 3713 to *H. macrotus*, omitting a series of characters that suggested an identity assignable to another taxon.

All the diagnostic characters of this specimen correspond to *H. diaphanopterus*, including strongly bicolored dorsal and ventral hairs (Fig. 3E), pale wing membranes, triangular translucent ears with a prominent lobe on the inner edge, and the presence of a gap between the ears with a membrane of approximately 4 mm height (Fig. 3F; Feijó et al. 2015).

Taxonomic remarks. Acosta and Venegas (2006) provided some external and cranial measurements of MNKM 3713, and they also reported a second specimen, MNKM 3714, of *H. macrotus*. However, Feijó et al. (2015) reidentified this second specimen as *H. diaphanopterus*. Given that the two previously referred individuals constitute the only records assigned to *H. macrotus* in Bolivia and that their identities do not correspond to this taxon, we exclude this species from the inventory of bat species known in Bolivia.

Family Emballonuridae

Cyttarops alecto Thomas, 1913

Material examined. BOLIVIA – Pando • Nicolás Suárez, Estación Biológica Tahuamanu; 11°20'11"S, 069°01'69"W; 287 m alt.; 06.IV.2009; Claudio J. Mamani leg.; 1 ♂ adult, CBG 401 (Figs. 2, 4A).

Identification. The specimen CBG 401 shows the following diagnostic traits described for this species: absence of glandular sacs in the uropatagium, thumb covered by the propatagium, skull with a cup-shaped depression, and narrow postorbital processes not fused

to the supraorbital ridges (Fig. 4A). Interestingly, CBG 401 and Peruvian representatives of *C. alecto* lack a diagnostic character previously reported for the species, namely the absence of a gap between the upper premolars (see Fig. 4A and Velazco et al. (2011) and Ludeña and Medina (2017) for Peruvian specimens). The Bolivian specimen shows values within the range of morphometric variation indicated for this species (Table 2).

Remarks. Aguirre et al. (2010) published the first record of this species for Bolivia, providing information on the geographic origin of the voucher specimen and the catalog number of the systematic collection where it was deposited (CBG 401). However, they did not include morphometric data or other information to support the identification of this specimen. The record reported here from Bolivia extends the distribution of *C. alecto* 770 km towards the southwestern Amazon (Fig. 2).

Saccopteryx canescens Thomas, 1901

Material examined. BOLIVIA – Santa Cruz • Parque Nacional Noel Kempff Mercado; 13°33'10"S, 061°00'51"W; 210 m alt.; 21.IX.1995; María C. Tapia leg.; 1 ♀ adult, MNKM 1910 (Figs. 2, 4B, 5A, B).

Saccopteryx leptura (Schreber, 1774)

Material examined. BOLIVIA – Santa Cruz • Parque Nacional Noel Kempff Mercado; 13°33'10"S, 061°00'51"W; 210 m alt.; 20.IX.1995; José L. Santivañez leg.; 1 ♀ adult, MNKM 1908 (Figs. 4C, 5C, D)

Identification. Most of the species of *Saccopteryx* Illiger, 1811 are characterized by having dorsal fur with two pale lines (except in *S. gymnura* Thomas, 1901 and *S. antioquiensis* Muñoz & Cuartas, 2001), with cranial traits that include a relatively flat rostrum, long and broad postorbital processes, and well-developed sagittal

Table 2. External and skull measurements (mm), in addition to weight (g), of representative specimens of *Cyttarops alecto*, *Saccopteryx canescens*, and *S. leptura* collected in Bolivia (CBG, MNKM) and other regions in South America. Range of measurements of *C. alecto* from Thomas (1913), Starrett and de la Torre (1964), Starrett and Casebeer (1968), Baker and Jones (1975), Masson and Cosson (1992), Ochoa et al. (1994), Velazco et al. (2011), Tavares et al. (2012), and Ludeña and Medina (2017). Range of measurements of *S. canescens* from Sanborn (1937), Carter and Dolan (1978), Husson (1962), Brosset and Charles-Dominique (1990), and Velazco et al. (2021).

	<i>Cyttarops alecto</i>			<i>S. canescens</i>			<i>S. leptura</i>
	Range	n	CBG 401	Range	n	MNKM 1910	MNKM 1908
Total length	70.00–90.00	4	70	—	—	61	63
Tail length	18.00–25.00	3	23	—	—	17	18
Hind foot length	8.00–11.20	5	8	4.80–7.30	7	7	8
Ear length	10.00–13.40	5	12	—	—	12	14
Forearm length	40.00–47.20	12	46.3	34.50–40.80	27	41.76	44.95
Body mass	5.60–9.00	5	—	—	—	4.50	6.00
Greatest length of skull	12.60–14.30	8	14.0	12.30–13.30	23	13.70	14.80
Condylar-canine length	12.26–12.70	3	13.0	11.20–11.60	4	12.26	13.40
Maxillary tooththrow length	5.10–5.70	10	5.6	4.60–5.10	23	5.30	5.60
Zygomatic breadth	8.00–8.88	8	8.9	7.70–8.50	22	—	9.29
Braincase breadth	6.70–7.10	8	7.1	6.10–6.50	5	6.85	7.18
Mastoid breadth	7.36–7.70	8	8.2	5.40–7.10	21	7.30	7.83
Postorbital breadth	3.70–4.50	9	4.0	2.10–3.20	21	2.65	2.13
Breadth across molars	5.86–6.30	5	6.4	5.10–5.80	22	5.54	6.44
Breadth across canines	2.40–3.35	6	2.9	2.60–3.00	21	3.10	3.34
Length of mandible	9.98	1	10.7	8.50–9.20	6	7.90	10.08
Mandibular tooththrow	—	—	5.7	5.00–6.00	7	5.30	5.81

crests. *Saccopteryx canescens* differs from its congeners by being small (forearm length 35.8–40.8 mm, upper dental row 4.6–5 mm, and width between molars 5.1–6 mm); the dorsal fur is gray or brown and lightly frosted, with two indistinct but evident dorsal lines (Hood and Gardner 2008). *Saccopteryx leptura* has dark brown dorsal fur and clearly visible dorsal lines, in addition to a larger size (forearm length 37.4–42.3 mm, upper dental row 5.1–5.5 mm) (Hood and Gardner 2008).

In comparison with an adult female of *S. leptura* (MNKM 1908; Figs. 4C, 5C, D) collected in the same locality, the specimen MNKM 1910 is smaller in external and cranial dimensions (Table 2), and the fur is more strongly bicolored (Fig. 5B), with dorsal and ventral hairs 8 and 6 mm long, respectively. Despite being smaller, the skull of *S. canescens* presents a greater postorbital width than *S. leptura*. Furthermore, the supraorbital ridges fuse well behind the postorbital constriction in *S. canescens*, whereas in *S. leptura* such fusion occurs almost at the level of the postorbital constriction (Table 2; Fig. 4B, C). All the chromatic characters of the fur and the body dimensions of *S. canescens* match the descriptions of this taxon, although some of its measurements exceed slightly the ranges indicated by other authors (Table 2).

Distribution. It is important to note that the geographic range of *S. canescens* as indicated by Hood and Gardner (2008), Solari (2015), Bonaccorso (2019), and Burgin et al. (2020) did not include Bolivia, even

though Emmons (1998) had recorded this species from the country. Fourteen specimens, including the one in this study, represent the southernmost records of this species; they are eight specimens from northwestern Brazil, five from Peru, and one from northeastern Bolivia (Sanborn 1937; Voss and Emmons 1996; Tejedor 2003; Hood and Gardner 2008; Velazco et al. 2021). The specimen recorded here (MNKM 1910) represents the southeastern most locality of *S. canescens*, extending the geographic range of the species 1133 km south of the closest previously known record in the southern extreme of Peru (Tejedor 2003; Fig. 2). The collection site of MNKM 1910 is an open savannah habitat on the edge of the Amazonian Forest, as previously reported for the species by Handley (1976), Jones and Hood (1993) and Emmons (1998).

Discussion

Our review of the existing data of bat species known to occur in Bolivia on the basis of voucher specimens has allowed us to generate a refined and updated checklist after the exclusion of one family and 13 species. Removed from the list of bat species in Bolivia are the following taxa (Table 1): in Emballonuridae, *Centronycteris maximiliani*, *Cormura brevirostris*, *Diclidurus albus*, and *Peropteryx kappleri*; in Phyllostomidae, *Anoura aequatoris*, *A. fistulata*, *A. peruana*, *Glyphoncteris behnii*, and *Micronycteris microtis*; in Furipteridae,



Figure 5. Dorsal and ventral views of the fur of *Saccopteryx canescens* and *S. leptura*. **A, B.** *S. canescens* (MNKM 1910). **C, D.** *S. leptura* (MNKM 1908).

Furipterus horrens (the only species of this family said to occur in Bolivia); in Molossidae, *Molossus currentium* and *M. pretiosus*; and in Vespertilionidae, *Histiotus macrotus*. Additional comments on these species, erroneously identified in previous lists of Bolivian bats, are below.

Centronycteris maximiliani—Venzal et al. (2013) reported a series of ectoparasites taken on nine specimens of this species, which was presumably collected in northeastern Bolivia (12°55'S, 062°52'W; 160 m alt.) near the Guapore River and on the border with Brazil. However, Venzal et al. (2013) did not provide information on the diagnostic traits of the referred specimens that may be used to verify the identity of the species. Furthermore, they did not mention the location of the referred specimens or if they were deposited in a systematic collection. Multiple attempts to contact the senior author on the paper (M.B. Labruna) were unsuccessful in securing access to the specimens. Therefore, it is impossible to confirm the identity of the bats assigned by Venzal et al. (2013) to *C. maximiliani* and, in fact, this species has not been included as part of the fauna of Bolivia in the most recent publications on the group (Sampaio et al. 2016; Bonaccorso 2019; Burgin et al. 2020).

Anoura spp.—According to recent revisionary work, eight South American species are recognized within the genus: *A. cadenai*, *A. caudifer* (including *A. aequatoris*), *A. cultrata*, *A. fistulata*, *A. geoffroyi* (including *A. peruana*), *A. javieri*, *A. latidens* (including *A. carishina*), and *A. luismanueli* (Calderón-Acevedo et al. 2021, 2022). In these publications, records from Bolivia said to be *A. aequatoris* (Mantilla-Meluk et al., 2012) belong instead to *A. caudifer*. Likewise, Bolivian records of *A. peruana* reported by Mantilla-Meluk and Baker (2010) are *A. geoffroyi* instead.

Micronycteris microtis—Recent studies using morphological and genetic data show that this species is a synonym *M. megalotis* (Table 1; Porter et al. 2007; Morales-Martínez et al. 2021), even though Díaz et al. (2021) and Simmons and Cirranello (2023) maintain *M. microtis* as a valid species. On the other hand, although the presence of *Molossus currentium* in the country is probable (Eger 2008; Barquez and Diaz 2016; Taylor et al. 2019; Burgin et al. 2020; Loureiro et al. 2020), its inclusion in the list of bats from Bolivia requires additional evidence.

Molossus pretiosus—This species was reported from Bolivia by Aguirre et al. (2010), based on three voucher specimens (CBG 402, 403, 404); however, these specimens could not be located in the CBG collection and are apparently lost. Aguirre et al. (2010) did not include information on external or cranial measurements, nor a discussion of the diagnostic characters used to identify these specimens. We temporarily recommend discarding *M. pretiosus* from the list of mammals of Bolivia, as suggested by other authors (Solari 2019; Taylor et al. 2019; Burgin et al. 2020), until the presence of this species can be corroborated by vouchered materials deposited in a systematic collection.

The new list of bat species confirmed from Bolivia contains eight families and 133 species (Table 3). The family Phyllostomidae represents the dominant fraction (76 species), followed by Vespertilionidae (21), Molossidae (20), Emballonuridae (8), Mormoopidae (3), Noctilionidae (2), Thyropteridae (2), and Natalidae (1). Two species are endemic to the country—*Micronycteris yatesi* Siles & Brooks, 2013 and *Eptesicus langeri* Acosta, Poma-Urey, Ossa-López, Rivera-Páez & Ramírez-Chaves, 2021—and eight are known by only one specimen—*Cyttarops alecto*, *Eptesicus brasiliensis* (Desmarest, 1819), *Gardnerycteris koepckeae* (Gardner & Patton, 1972), *Lichonycteris degener* Miller, 1931, *Lionycteris spurrelli* Thomas, 1913, *Lophostoma carrikeri* (J.A. Allen, 1910), *Peropteryx pallidoptera* Lim, Engstrom, Reid, Simmons, Voss & Fleck, 2010, and *Saccopteryx canescens*.

Several bat species are known to occur in Bolivia by very few records, just one or two specimens, and have not been reported in the last 30 years. In this group we highlight *Eumops hansae* (only two individuals collected in 1986 and 1992; Ibáñez and Ochoa 1989; Aguirre and Urioste 1994; Anderson 1997), *Lichonycteris degener* (collected in 1979; Anderson et al. 1982), and *Lophostoma carrikeri* (collected in 1964; Koopman 1976). Therefore, we highlight the need to assess the conservation status of these underreported, rare species of bats. Furthermore, exhaustive fieldwork is needed to sample poorly known areas, where there is the greatest chances of success in surveying the fauna. These initiatives must consider the use of conventional and unconventional sampling methods, including ultrasonic detection (Schnitzler and Kalko 2001; Jung et al. 2014) accompanied by acoustic lures in some cases (Hintze et al. 2020), principally for the inventory of aerial insectivorous bats. The reliability of the data provided by this technology has increased to the extent that species' echolocation patterns have been described and are easily distinguishable, but it is essential that these data are accompanied by reference images of typical search phase calls or the descriptions of fundamental parameters (Ochoa et al. 2000; Jung et al. 2007, 2014; Corben and Miller pers. comm.). The availability of acoustic files in libraries or on web pages, such as the ChiroVox bat call database (Görföl 2022), are an important tool for the recognition and the comparison of echolocation patterns as a complement to the collection of duly identified voucher specimens in cases of little-known or cryptic species (e.g., *Centronycteris centralis* Thomas, 1912; Espinal et al. 2021).

Conventional methods to detect or collect bats must include sampling efforts with mist nets in various strata, including the canopy level, and the systematic review of roosts, in addition to the eventual capture of aerial insectivorous bats with shotguns and spotlights in open spaces. All these alternatives, supported by ethical protocols, have demonstrated excellent results in the Neotropics (e.g., Handley 1976; Simmons and Voss 1998; Ochoa et al. 2005, 2008, 2009; Díaz and




Figure 6. Species-accumulation curve of bat surveys in Bolivia during the period 1885–2016, showing the importance of the expeditions conducted between 1963 and 1981 by the Middle America Research Unit, AMNH, LSUMZ, the Zoológico de Santa Cruz and EBD, the Museum of Southwestern Biology, and the CBF (see Table 3).

Linares 2012; Jung et al. 2014). Finally, it is extremely important that researchers of future faunal surveys must assume the implicit responsibility in careful taxonomic identification, taking into account the comparative analyses of voucher specimens, consultation of the specialized literature, and the advice of local or regional experts.

Despite the reduction in 8.9% of the species richness of bats of Bolivia as a result of our study (eight families and 133 species versus nine families and 146 species according to Díaz et al. 2021), the country harbors a high diversity of bats with respect to other Neotropical countries. The collection of voucher specimens represents a fundamental process to preserve key evidence of the distributions and taxonomic identifications of mammal species (López-Vidal and Elizalde-Arellano 2006). In Bolivia, taxonomic inventories of bats have been mostly based on the comparative study of preserved specimens, which have enriched the existing biological heritage in systematic collections (Solari and Baker 2006; Moratelli and Wilson 2014; Velazco and Patterson 2019; Poma-Urey et al. 2021).

Among the taxa that make up the list of bats in Bolivia, 44.8% (60 species) were recorded for the first time between 1960 and 1986, with an important additional contribution in the last 36 years; however, the species-accumulation curve still shows an increasing trend that suggests the need for greater inventory efforts (Fig. 6). The knowledge accumulated in the last decades includes the invaluable fieldwork of at least 17 national and approximately 40 foreign bat researchers who have generated the data supporting this study (Table 3). Among them, we highlight the works of S. Anderson, J. Steinbach, A. Ximénez, W.D. Webster, C. Ibáñez, C.C. Olrog, C.G. Schmitt, G.K. Creighton, K.F. Koopman, M.L. Kuns, L.H. Acosta, L. Aguirre, D.E.

Añez, H. Azurduy, M. Hidalgo, F. Hinojosa, L. Maffei, A. Muñoz, J. Salazar-Bravo, J.L. Santivañez, L. Siles, M.C. Tapia-Arauz, and A. Vargas.

Acknowledgements

We thank R. Barquez (Colección Mamíferos Lillo), L.F. Aguirre and F. Navarro (Centro de Biodiversidad y Genética), A. Rico (Colección Boliviana de Fauna), P. Cordeiro (Universidade Federal da Paraíba), C. Ibáñez (Estación Biológica Doñana) and L.F. Silveira (Museu de Zoologia da Universidade de São Paulo), for allowing us to study specimens under their care and for all their contributions during this research. L. Loureiro and B. Lim provided data on *Molossus fluminensis*, M. Herrera helped to locate the MZUSP specimen of *Peropteryx macrotis*, and M. Terán provided data from the second specimen of *Micronycteris hirsuta* recorded for Bolivia. The reviewers made valuable suggestions for improving previous versions of this document.

References

- Acosta L, Aguanta F (2005) Nota sobre un nuevo registro de murciélago (*Lampronnycteris brachyotis*) para Bolivia. *Kempffiana* 1: 65–69.
- Acosta L, Venegas C (2006) Algunas consideraciones taxonómicas de *Histiotus laephotis* e *H. macrotus*, en Bolivia. *Kempffiana* 2 (1): 109–115.
- Acosta L, Azurduy H, Sánchez G, Cortez E (2006) Primer registro *Neoplatymops mattogrossensis* (Vieira, 1942) (Chiroptera), en Bolivia. *Kempffiana* 2 (1): 134–139.
- Acosta LH, Poma-Urey JL, Ossa-López PA, Rivera-Páez FA, Ramírez-Chaves HE (2021) A new species of *Eptesicus* (Mammalia: Chiroptera: Vespertilionidae), from the sub-Andean Forest of Santa Cruz, Bolivia. *Therya* 12 (3): 391–409. <https://doi.org/10.12933/therya-21-1119>

Table 3. Check list of the bats with presence in Bolivia supported by voucher specimens. References correspond to the first published and documented records for the country.

Taxa	Voucher	Collector(s)	Year	Department	Reference(s)
Family Emballonuridae (8)					
<i>Cyttarops alecto</i> Thomas, 1913	CBG 401	C. Mamani	2009	Pando	Aguirre et al. 2010; this study
<i>Peropteryx leucoptera</i> Peters, 1867	AMNH 209228	K.F. Koopman	1964	Beni	Poma-Urey et al. 2021
<i>Peropteryx macrotis</i> (Wagner, 1843)	CML 1325	C.C. Olrog	1959	Santa Cruz	Barquez 1983
<i>Peropteryx pallidoptera</i> Lim, Engstrom, Reid, Simmons, Voss & Fleck, 2010	MHNC-M 555	M. Hidalgo	2016	Cochabamba	Siles et al. 2021
<i>Rhynchonycteris naso</i> (Wied-Neuwied, 1820)	AMNH 209187	S. Anderson	1964	Beni	Anderson et al. 1982
<i>Saccopteryx bilineata</i> (Temminck, 1838)	FMNH 21474	J. Steinbach	1914	Santa Cruz	Sanborn 1937
<i>Saccopteryx canescens</i> Thomas, 1901	MNKM 1910	M.C. Tapia-Arauz	1995	Santa Cruz	Emmons 1998; this study
<i>Saccopteryx leptura</i> (Schreber, 1774)	AMNH 209225	S. Anderson	1964	Beni	Anderson et al. 1982
Family Phyllostomidae (76)					
Subfamily Carollinae					
<i>Carollia benkeithi</i> Solari & Baker, 2006	AMNH 246548	D.C. Cole	1979	La Paz	Solari and Baker 2006
<i>Carollia brevicauda</i> (Schinz, 1821)	FMNH 21482	J. Steinbach	1914	Santa Cruz	Pine 1972
<i>Carollia manu</i> Pacheco, Solari & Velazco, 2004	AMNH 264995	C.T. Seaton	1993	La Paz	Pacheco et al. 2004
<i>Carollia perspicillata</i> (Linnaeus, 1758)	AMNH 61756	J. Steinbach	1921	Santa Cruz	Anderson et al. 1982
Subfamily Desmodontinae					
<i>Desmodus rotundus</i> (É. Geoffroy, 1810)	BMNH 2.1.1.3	P.O. Simons	1901	Cochabamba	Thomas 1902
<i>Diaemus youngii</i> (Jentink, 1893)	AMNH 209743	S. Anderson	1964	Beni	Koopman 1976
<i>Diphylla ecaudata</i> Spix, 1823	AMNH 261777	L.A. Ruedas	1985	La Paz	Anderson 1991
Subfamily Glossophaginae					
<i>Anoura caudifer</i> (É. Geoffroy, 1818)	AMNH 246468	C.G. Schmitt	1979	La Paz	Anderson et al. 1982
<i>Anoura cultrata</i> Handley, 1960	LSUMZ 22962	S.W. Cardiff	1980	La Paz	Anderson et al. 1982
<i>Anoura geoffroyi</i> Gray, 1838	FMNH 39294	J.D. Haseman	1909	Santa Cruz	Sanborn 1933
<i>Anoura latidens</i> Handley, 1984	AMNH 264604	S. Anderson	1992	La Paz	Calderón-Acevedo and Muchhala 2020
<i>Choeroniscus minor</i> (Peters, 1868)	TTU-M 34789	W.D. Webster	1979	La Paz	Webster and Jones 1980
<i>Glossophaga soricina</i> (Pallas, 1766)	AMNH 61765	J. Steinbach	1921	Santa Cruz	Anderson et al. 1982
<i>Lichonycteris degener</i> Miller, 1931	AMNH 244621	S. Anderson	1979	Cochabamba	Anderson et al. 1982
Subfamily Glyphonycterinae					
<i>Glyphonycteris daviesi</i> (Hill, 1964)	CBF 2908	K.S. Smith	1993	La Paz	Anderson 1997
<i>Trinycteris nicefori</i> (Sanborn, 1949)	CBF 2892	F. Hinojosa	1990	La Paz	Emmons 1991
Subfamily Lonchophyllinae					
<i>Hsunnycteris pattoni</i> (Woodman & Timm, 2006)	AMNH 209358	D.E. Añez	1964	Beni	Parlos et al. 2014
<i>Hsunnycteris thomasi</i> (J.A. Allen, 1904)	AMNH 262429	C.K. Malcolm	1986	Pando	Anderson 1997
<i>Lionycteris spurrelli</i> Thomas, 1913	MNKM 3283	H. Azurduy	2002	Santa Cruz	Azurduy and Emmons 2005
<i>Lonchophylla dekeyseri</i> Taddei, Vizotto & Sazima, 1983	USNM 584472	L.H. Emmons	2000	Santa Cruz	Moratelli and Dias 2015
Subfamily Lonchorhininae					
<i>Lonchorhina aurita</i> Tomes, 1863	CM 1890	J.D. Haseman	1909	Santa Cruz	Sanborn 1932a
Subfamily Micronycterinae					
<i>Lampronnycteris brachyotis</i> (Dobson, 1879)	MNKM 3730	L.H. Acosta	2003	Santa Cruz	Acosta and Aguanta 2005
<i>Micronycteris hirsuta</i> (Peters, 1869)	MNKM 3267	H. Azurduy	2002	Santa Cruz	Azurduy and Emmons 2005
<i>Micronycteris megalotis</i> (Gray, 1842)	UMMZ 126843	G.K. Creighton	1978	Cochabamba	Anderson et al. 1982
<i>Micronycteris minuta</i> (Gervais, 1856)	UMMZ 126729	G.K. Creighton	1978	La Paz	Anderson et al. 1982
<i>Micronycteris sanborni</i> Simmons, 1996	MNKM 1988	L. Maffei	1996	Santa Cruz	Poma-Urey et al. 2020
<i>Micronycteris yatesi</i> Siles & Brooks, 2013	MHNC-M 157	L. Siles, A. Muñoz	2007	Chuquisaca	Siles et al. 2013

Taxa	Voucher	Collector(s)	Year	Department	Reference(s)
Subfamily Phyllostominae					
<i>Chrotopterus auritus</i> (Peters, 1856)	AMNH 209353	S. Anderson	1964	Beni	Anderson et al. 1982
<i>Gardnerycteris crenulatum</i> (É. Geoffroy, 1803)	AMNH 209323	S. Anderson	1964	Beni	Koopman 1976
<i>Gardnerycteris koepckeae</i> (Gardner & Patton, 1972)	CBF 8683	L. Siles	2016	La Paz	Siles and Wallace 2021
<i>Lophostoma brasiliense</i> Peters, 1866	CBF 179	J. Salazar-Bravo	1987	Beni	Wilson and Salazar-Bravo 1990
<i>Lophostoma carrikeri</i> (J.A. Allen, 1910)	AMNH 209322	S. Anderson	1964	Beni	Koopman 1976
<i>Lophostoma silvicolum</i> d'Orbigny, 1836	FMNH 44867	J. Steinbach	1911	Santa Cruz	Davis and Carter 1978
<i>Macrophyllum macrophyllum</i> (Schinz, 1821)	AMNH 209320	S. Anderson	1964	Beni	Harrison 1975
<i>Phylloderma stenops</i> Peters, 1865	CML 413	R. Ojeda, R.M. Barquez	1975	Santa Cruz	Barquez and Ojeda 1979
<i>Phyllostomus discolor</i> (Wagner, 1843)	AMNH 209325	A. Ximénez	1964	Beni	Anderson et al. 1982
<i>Phyllostomus elongatus</i> (É. Geoffroy, 1810)	AMNH 209331	K.F. Koopman	1964	Beni	Anderson et al. 1982
<i>Phyllostomus hastatus</i> (Pallas, 1767)	AMNH 61751	J. Steinbach	1921	Santa Cruz	Anderson et al. 1982
<i>Tonatia bidens</i> (Spix, 1823)	MNKM 4745	F. Aguanta	2005	Santa Cruz	Paca et al. 2012
<i>Tonatia maresi</i> Williams, Willig & Reid, 1995	CBF 2849	L.H. Emmons	1992	Pando	Emmons and Smith 2002
<i>Trachops cirrhosus</i> (Spix, 1823)	AMNH 209348	A. Ximénez	1964	Beni	Koopman 1976
<i>Vampyrum spectrum</i> (Linnaeus, 1758)	AMNH 261379	J.A. Cook	1985	Beni	Anderson 1991
Subfamily Rhinophyllinae					
<i>Rhinophylla pumilio</i> Peters, 1865	MSU 32895	W.D. Webster	1979	La Paz	Webster and Jones 1980
Subfamily Stenodermatinae					
<i>Artibeus lituratus</i> (Olfers, 1818)	AMNH 1255	H.H. Rusby	1885	Cochabamba	Allen 1904
<i>Artibeus obscurus</i> (Schinz, 1821)	AMNH 38502	L.E. Miller, H.S. Boyle	1915	Cochabamba	Anderson 1997
<i>Artibeus planirostris</i> (Spix, 1823)	TTU-M 34863	W.D. Webster	1979	La Paz	Webster & Jones 1980
<i>Artibeus (Koopmania) concolor</i> Peters, 1865	CBG 450	A. Vargas	2006	Pando	Vargas and Balderrama 2009
<i>Dermanura anderseni</i> (Osgood, 1916)	CM 2734	J. Steinbach	1911	Santa Cruz	Sanborn 1932a
<i>Dermanura glauca</i> (Thomas, 1893)	CML 2207	C.C. Olrog	1976	Cochabamba	Barquez and Olrog 1980
<i>Dermanura gnoma</i> (Handley, 1987)	LSUMZ 32966	W.D. Webster	1979	La Paz	Webster and Jones 1980
<i>Chiroderma salvini</i> Dobson, 1878	AMNH 246625	C.G. Schmitt	1979	La Paz	Anderson et al. 1982
<i>Chiroderma trinitatum</i> Goodwin, 1958	AMNH 209520	K.F. Koopman	1964	Beni	Koopman 1976
<i>Chiroderma villosum</i> Peters, 1860	AMNH 61754	J. Steinbach	1921	Santa Cruz	Koopman 1976
<i>Enchisthenes hartii</i> (Thomas, 1901)	TTU-M 34880	W.D. Webster	1979	La Paz	Webster and Jones 1980
<i>Mesophylla macconnelli</i> Thomas, 1901	AMNH 209577	R.G. Van Gelder	1964	Beni	Koopman 1976
<i>Platyrrhinus albericoi</i> Velazco, 2005	CML 1581	C.C. Olrog	1976	Cochabamba	Barquez and Olrog 1980
<i>Platyrrhinus brachycephalus</i> (Rouk & Carter, 1972)	UMMZ 126755	G.K. Creighton	1978	Cochabamba	Velazco 2005
<i>Platyrrhinus incarum</i> (Thomas, 1912)	USNM 461044	V. Adcock	1971	Cochabamba	Anderson 1997
<i>Platyrrhinus infuscus</i> (Peters, 1880)	CML 2215	C.C. Olrog	1976	Cochabamba	Barquez and Olrog 1980
<i>Platyrrhinus lineatus</i> (É. Geoffroy, 1810)	AMNH 210805	A. Ximénez	1965	Beni	Anderson 1997
<i>Platyrrhinus masu</i> Velazco, 2005	AMNH 246610	C.G. Schmitt	1979	La Paz	Velazco 2005
<i>Platyrrhinus umbratus</i> (Lyon, 1902)	UMMZ 127174	G.K. Creighton	1978	La Paz	Velazco et al. 2018
<i>Pygoderma bilabiatum</i> (Wagner, 1843)	AMNH 246399	R.M. Barquez, R. Ojeda	1975	Santa Cruz	Ojeda and Barquez 1978
<i>Sphaeronycteris toxophyllum</i> Peters, 1882	AMNH 209740	S. Anderson	1964	Beni	Koopman 1976
<i>Sturnira erythromos</i> (Tschudi, 1844)	MSU 32905	W.D. Webster	1979	La Paz	Webster and Jones 1980
<i>Sturnira giannae</i> Velazco & Patterson, 2019	AMNH 209418	A. Ximénez	1964	Beni	Velazco and Patterson 2019
<i>Sturnira liliium</i> (É. Geoffroy, 1810)	CM 1944	J. Steinbach	1909	Santa Cruz	Sanborn 1932a
<i>Sturnira magna</i> de la Torre, 1966	TTU-M 34907	W.D. Webster	1979	La Paz	Webster and Jones 1980
<i>Sturnira oporaphilum</i> (Tschudi, 1844)	LSUMZ 23065	T.A. Parker	1979	La Paz	Webster and Jones 1980

Taxa	Voucher	Collector(s)	Year	Department	Reference(s)
<i>Sturnira sorianoi</i> Sánchez-Hernández, Romero-Almaraz & Schnell, 2005.	AMNH 260870	C.G. Schmitt	1984	Santa Cruz	Sánchez-Hernández et al. 2005
<i>Sturnira tildae</i> de la Torre, 1959	AMNH 209419	A. Ximénez	1964	Beni	Anderson et al. 1982
<i>Uroderma bilobatum</i> Peters, 1866	BMNH 1.2.1.37	P.O. Simons	1900	Beni	Davis 1968
<i>Uroderma magnirostrum</i> Davis, 1968	AMNH 209425	S. Anderson	1964	Beni	Davis 1968
<i>Vampyressa thylene</i> Thomas, 1909	AMNH 262524	T.L. Yates	1986	Pando	Anderson 1991
<i>Vampyriscus bidens</i> (Dobson, 1878)	LSUMZ 26652	C.M. Fugler	1981	Beni	Anderson and Webster 1983
<i>Vampyrodes caraccioli</i> (Thomas, 1889)	AMNH 209518	K.F. Koopman	1964	Beni	Anderson et al. 1982
Mormoopidae (3)					
<i>Pteronotus gymnonotus</i> (Wagner, 1843)	EBD 14269	C. Ibáñez	1986	Santa Cruz	Ibáñez and Ochoa 1989
<i>Pteronotus personatus</i> (Wagner, 1843)	EBD 14294	C. Ibáñez	1986	Santa Cruz	Ibáñez and Ochoa 1989
<i>Pteronotus rubiginosus</i> (Wagner, 1843)	EBD 14267	C. Ibáñez	1986	Santa Cruz	Ibáñez and Ochoa 1989
Noctilionidae (2)					
<i>Noctilio albigentris</i> Dermarest, 1818	AMNH 210540	S. Anderson	1965	Beni	Anderson et al. 1982
<i>Noctilio leporinus</i> (Linnaeus, 1758)	AMNH 210666	A. Ximénez	1965	Beni	Anderson et al. 1982
Thyropteridae (2)					
<i>Thyroptera discifera</i> (Lichtenstein & Peters, 1855)	MNKM 1907	J.L. Santivañez	1995	Santa Cruz	Emmons et al. 2006
<i>Thyroptera tricolor</i> Spix, 1823	LSUMZ 26686	C.M. Fugler	1981	Beni	Anderson and Webster 1983
Natalidae (1)					
<i>Natalus macrourus</i> (Gervais, 1856)	SJRP 14507	W. Uieda, A. Rodrigues	1983	Santa Cruz	Taddei and Uieda 2001
Molossidae (20)					
<i>Cynomops abrasus</i> (Temminck, 1826)	EBD 13520	C. Ibáñez, F. Braza	1983	Santa Cruz	Ibáñez 1985
<i>Cynomops planirostris</i> (Peters, 1866)	FMNH 96038	M.L. Kuns	1963	Beni	Anderson et al. 1982
<i>Eumops auripendulus</i> (Shaw, 1800)	USNM 239009	W.M. Mann	1921	La Paz	Sanborn 1932b
<i>Eumops chimaera</i> Gregorin, Moras, Acosta, Vasconcellos, Poma, dos Santos & Paca, 2016.	MNKM 4753	R.C. Paca	2009	Santa Cruz	Gregorin et al. 2016
<i>Eumops glaucinus</i> (Wagner, 1843)	USNM 390643	G.L. Ranck	1966	Beni	Eger 1977
<i>Eumops hansae</i> Sanborn, 1932	EBD 14295	C. Ibáñez	1986	Santa Cruz	Ibáñez and Ochoa 1989
<i>Eumops patagonicus</i> Thomas, 1924	FMNH 21536	R.H. Becker	1915	Beni	Sanborn 1932b
<i>Eumops perotis</i> (Schinz, 1821)	FMNH 116690	M.L. Kuns	1963	Beni	Anderson and Webster 1983
<i>Eumops trumbulli</i> (Thomas, 1901)	AMNH 209900	A. Ximénez	1964	Beni	Eger 1977
<i>Molossops temminckii</i> (Burmeister, 1854)	FMNH 115814	M.L. Kuns	1963	Beni	Anderson et al. 1982
<i>Molossus fluminensis</i> Lataste, 1891	AMNH 263285	S. Anderson	1987	Santa Cruz	Loureiro et al. 2020
<i>Molossus molossus</i> (Pallas, 1766)	CM 1892	J.D. Haseman	1909	Beni	Anderson 1991
<i>Molossus rufus</i> É. Geoffroy, 1805	AMNH 211279	D.E. Añez	1965	Beni	Anderson et al. 1982
<i>Neoplattymops mattogrossensis</i> (Vieira, 1942)	MNKM 3710	G. Sánchez, E. Cortez	2006	Santa Cruz	Acosta et al. 2006
<i>Nyctinomops aurispinosus</i> (Peale, 1848)	EBD 14282	C. Ibáñez	1986	Santa Cruz	Ibáñez and Ochoa 1989
<i>Nyctinomops laticaudatus</i> (É. Geoffroy, 1805)	AMNH 209774	S. Anderson	1964	Beni	Anderson et al. 1982
<i>Nyctinomops macrotis</i> (Gray, 1840)	EBD 14046	C. Ibáñez	1986	Santa Cruz	Ibáñez and Ochoa 1989
<i>Promops centralis</i> Thomas, 1915	AMNH 260273	S. Anderson	1984	Santa Cruz	Anderson 1991
<i>Promops nasutus</i> (Spix, 1823)	AMNH 260306	S. Anderson	1984	Santa Cruz	Anderson 1997
<i>Tadarida brasiliensis</i> (L. Geoffroy, 1824)	CM 5278	J. Steinbach	1921	Cochabamba	Sanborn 1932a
Vespertilionidae (21)					
<i>Eptesicus andinus</i> J.A. Allen, 1914	BMNH 2.1.1.1	M.R. O. Thomas	1901	Cochabamba	Acosta et al. 2021
<i>Eptesicus brasiliensis</i> (Dermarest, 1819)	AMNH 260257	N. Olds	1984	Santa Cruz	Poma-Urey et al. 2019
<i>Eptesicus chiriquinus</i> Thomas, 1920	MHNC-M 154	L. Siles	2007	Chuquisaca	Siles 2007
<i>Eptesicus diminutus</i> Osgood, 1915	MNKM 5658	A.M. Osinaga	1997	Santa Cruz	Poma-Urey et al. 2019

Taxa	Voucher	Collector(s)	Year	Department	Reference(s)
<i>Eptesicus furinalis</i> (d'Orbigny & Gervais, 1847)	LSUMZ 26690	C.M. Fugler	1981	Beni	Anderson et al. 1982
<i>Eptesicus langeri</i> Acosta, Poma-Urey, Ossa-López, Rivera-Páez & Ramírez-Chaves, 2021	MNKM 5584	L.H. Acosta	2013	Santa Cruz	Acosta et al. 2021
<i>Histiotus diaphanopterus</i> Feijó, da Rocha & Althoff, 2015	AMNH 264086	J. Salazar-Bravo	1991	Santa Cruz	Semedo and Feijó 2016
<i>Histiotus laephotis</i> Thomas, 1916	BMNH 97.2.25.1	A. Borelli	1896	Tarija	Thomas 1916
<i>Histiotus montanus</i> (Philippi & Landbeck, 1861)	EBD 13256	J. Cabot-Nieves, P. Serrano	1983	La Paz	Ibáñez 1985
<i>Histiotus velatus</i> (L. Geoffroy, 1824)	CBF 206	F. Hinojosa	1989	La Paz	Anderson 1997
<i>Lasiurus blossevillii</i> (Lesson & Garnot, 1826)	AMNH 209773	A. Ximénez	1964	Beni	Anderson et al. 1982
<i>Lasiurus ega</i> (Gervais, 1856)	FMNH 21476	J. Steinbach	1915	Santa Cruz	Anderson et al. 1982
<i>Lasiurus villosissimus</i> É. Geoffroy, 1806	LSUMZ 22703	J.V. Jr. Remsen	1980	La Paz	Anderson et al. 1982
<i>Myotis albescens</i> (É. Geoffroy, 1806)	AMNH 209767	S. Anderson	1964	Beni	LaVal 1973
<i>Myotis dinellii</i> Thomas, 1902	AMNH 39003	L.E. Miller, H.S. Boyle	1915	Potosí	Anderson 1997
<i>Myotis keaysi</i> J.A. Allen, 1914	AMNH 23161	J.V. Jr. Remsen	1979	La Paz	Anderson et al. 1982
<i>Myotis midastactus</i> Moratelli & Wilson, 2014	AMNH 211156	S. Anderson	1965	Beni	Moratelli and Wilson 2014
<i>Myotis nigricans</i> (Schinz, 1821)	AMNH 38503	L.E. Miller, H.S. Boyle	1915	Cochabamba	Anderson et al. 1982
<i>Myotis oxyotus</i> (Peters, 1866)	BMNH 2.2.2.125	P.O. Simons	1902	La Paz	LaVal 1973
<i>Myotis riparius</i> Handley, 1960	AMNH 260251	N. Olds	1984	Santa Cruz	Anderson 1991
<i>Rhogeessa io</i> Thomas, 1903	FMNH 115807	M.L. Kuns	1963	Beni	Anderson et al. 1982

- Aguirre LF** (2007) Historia natural, distribución y conservación de los murciélagos de Bolivia. Fundación Simón I. Patiño. Santa Cruz, Bolivia, 416 pp.
- Aguirre LF, Urioste RJ** (1994) Nuevos registros de murciélagos para Bolivia y los departamentos de Beni y Pando. *Ecología en Bolivia* 23: 71–76.
- Aguirre LF, Mamani CJ, Barboza-Márquez K, Mantilla-Meluk H** (2010) Lista actualizada de los murciélagos de Bolivia. *Revista Boliviana de Ecología y Conservación Ambiental* 27: 1–7.
- Aguirre LF, Tarifa T, Wallace RB, Bernal H, Siles L, Aliaga-Rossel E, Salazar-Bravo J** (2019) Lista actualizada y comentada de los mamíferos de Bolivia. *Ecología en Bolivia* 54 (2): 108–148.
- Alfaro-Lara H, Villalobos-Chaves D, Castillo-Gómez JJ, Zuñiga-Montero AF, Pineda-Lizano W** (2018) Rediscovery of *Furipterus horrens* (Chiroptera: Furipteridae) in Costa Rica. *Mammalia* 83 (4): 352–356. <https://doi.org/10.1515/mammalia-2018-0066>
- Allen JA** (1904) New bats from tropical America, with notes on species of *Otopterus*. *Bulletin American Museum of Natural History* 20: 227–237.
- Anderson S, Koopman KF, Creighton GK** (1982) Bats of Bolivia: an annotated checklist. *Bulletin of the American Museum of Natural History* 2750: 1–24.
- Anderson S, Webster WD** (1983) Notes on Bolivian mammals. 1. Additional records of bats. *American Museum Novitates* 2766: 1–3.
- Anderson S** (1991) A brief-history of Bolivian chiroptology and new records of bats. *Bulletin of the American Museum of Natural History* 206: 138–144.
- Anderson S** (1997) Mammals of Bolivia: taxonomy and distribution. *Bulletin of the American Museum of Natural History* 231: 1–652.
- Arroyo-Cabrales J** (2019) Family Furipteridae. In: Wilson DE, Mittermeier RA (Eds.) *Handbook of the mammals of the world, bats, volume 9*. Lynx Editions. Barcelona, Spain, 412–417.
- Azurduy H, Emmons LH** (2005) Nota sobre dos nuevas especies de murciélagos para Bolivia. *Ecología en Bolivia* 40 (2): 53–57.
- Baker RJ, Jones JK Jr** (1975) Additional records of bats from Nicaragua, with a revised checklist of Chiroptera. *Occasional Papers (The Museum, Texas Tech University)* 32: 1–13. <https://doi.org/10.5962/bhl.title.142873>
- Barquez** (1983) Una nueva localidad para la distribución de *Peropteryx macrotis macrotis* (Wagner) (Chiroptera, Emballonuridae). *Historia Natural* 3 (21): 185–186.
- Barquez R, Díaz M** (2016) *Molossus currentium*. The IUCN Red List of Threatened Species 2016: e.T88087340A22107231. <https://doi.org/10.2305/iucn.uk.2016-3.rlts.t88087340a22107231.en>. Accessed on: 2022-11-10.
- Barquez RM, Ojeda RA** (1979) Nueva subespecie de *Phylloderma stenops* (Chiroptera Phyllostomidae). *Neotropica* 25: 83–89.
- Barquez RM, Olrog CC** (1980) Tres nuevas especies de *Vampyrops* para Bolivia (Chiroptera Phyllostomidae). *Neotropica* 26: 53–56.
- Barquez RM, Mares MA, Braun JK** (1999) The bats of Argentina. *Special Publications of the Museum of Texas Tech University* 42: 1–275.
- Basantes M, Tinoco N, Velazco PM, Hofmann MJ, Rodríguez-Posada ME, Camacho MA** (2020) Systematics and Taxonomy of *Tonatia saurophila* Koopman &

- Williams, 1951 (Chiroptera, Phyllostomidae). *ZooKeys* 915: 59–86. <https://doi.org/10.3897/zookeys.915.46995>
- Bonaccorso F** (2019) Family Emballonuridae. In: Wilson DE, Mittermeier RA (Eds.) *Handbook of the mammals of the world, bats, volume 9*. Lynx Edicions. Barcelona, Spain, 334–373.
- Brosset A, Charles-Dominique P** (1990) The bats from French Guiana: a taxonomic, faunistic and ecological approach. *Mammalia* 54: 509–560.
- Burgin CJ, Wilson DE, Mittermeier RA, Rylands AB, Lacher TE, Secrest W** (2020) *Illustrated checklist of mammals of the world, Monotremata to Rodentia, volume 1*. Lynx Edicions. Barcelona, Spain, 1166 pp.
- Cabot-Nieves J** (1996) La expedición zoológica hispano-boliviana al Parque Nacional de Huanchaca. Landívar SRL. Santa Cruz, Bolivia, 56 pp.
- Calderón-Acevedo CA, Muchhala NC** (2018) Identification and diagnosis of *Anoura fistulata* with remarks on its presumed presence in Bolivia. *Journal of Mammalogy* 99 (1): 131–137. <https://doi.org/10.1093/jmammal/gyx159>
- Calderón-Acevedo CA, Muchhala N** (2020) First report of the Broad-toothed Tailless Bat, *Anoura latidens* Handley, 1984 (Chiroptera, Phyllostomidae), in Bolivia. *Check List* 16 (6): 1545–1550. <https://doi.org/10.15560/16.6.1545>
- Calderón-Acevedo CA., Rodríguez-Posada ME, Muchhala N** (2021) Morphology and genetics concur that *Anoura carishina* is a synonym of *Anoura latidens* (Chiroptera, Glossophaginae). *Mammalia* 85 (5): 471–481. <https://doi.org/10.1515/mammalia-2020-0183>
- Calderón-Acevedo CA, Bagley JC, Muchhala N** (2022) Genome-wide ultraconserved elements resolve phylogenetic relationships and biogeographic history among Neotropical leaf-nosed bats in the genus *Anoura* (Phyllostomidae). *Molecular Phylogenetics and Evolution* 167: 107356. <https://doi.org/10.1016/j.ympev.2021.107356>
- Carter DC, Dolan PG** (1978) *Catalogue of type specimens of neotropical bats in selected European museums*. Special Publications of the Museum 15. Texas Tech University Press, Lubbock, USA, 136 pp.
- Davis WB** (1968) Review of the genus *Uroderma* (Chiroptera). *Journal of Mammalogy* 49 (4): 676–698. <https://doi.org/10.2307/1378728>
- Davis WB, Carter D** (1978) A review of the round-eared bats of the *Tonatia silvicola* complex, with descriptions of three new taxa. *Occasional Papers Museum of Texas Tech University* 53: 1–12.
- Díaz M, Linares VH** (2012) Refugios naturales y artificiales de murciélagos (Mammalia: Chiroptera) en la selva baja en el noroeste de Perú. *Gayana* 76 (2): 117–130.
- Díaz M, Solari S, Gregorin R, Aguirre LF, Barquez R** (2021) Clave de identificación de los murciélagos neotropicales/Chave de identificação dos morcegos Neotropicais. Bilingüe: español-portugués. *Publicación Especial N° 2. Programa de Conservación de los Murciélagos de Argentina, Tucumán, Argentina*, 207 pp.
- Eger JL** (1977) Systematics of the genus *Eumops* (Chiroptera: Molossidae). *Life Sciences Contributions, Royal Ontario Museum* 110: 1–69.
- Eger JL** (2008) Family Molossidae Gervais, 1856. In: Gardner AL (Ed.) *Mammals of South America, vol. 1, marsupials, xenarthrans, shrews, and bats*. University of Chicago Press, Chicago, USA, 399–440.
- Emmons LH** (1991) Mammal of Alto Madidi In: Parker T, Bailey B (Eds.) *A biological assessment of the Alto Madidi Region and adjacent areas of northwest Bolivia*. RAP Working Papers 1: 72–73.
- Emmons LH** (1998) Mammal fauna of Parque Noel Kempff Mercado. In: Killeen TJ, Schulenberg TS (Eds.) *A biological assessment of Parque Nacional Noel Kempff Mercado, Bolivia*. RAP Working Papers 10: 129–143.
- Emmons LH, Smith KS** (2002) Mamíferos del noroeste de la zona de Pando. In: Montambault RJ (Ed.) *Informes de las evaluaciones biológicas Pampas del Heath, Perú, Alto Madidi, Bolivia y Pando*. RAP Working Papers 24: 111–113.
- Emmons LH, Swarner MJ, Vargas-Espinoza A, Tschapka M, Azurduy H, Kalko EKV** (2006) The forest and savanna bat communities of Noel Kempff Mercado National Park (Bolivia). *Revista Boliviana de Ecología y Conservación Ambiental* 19: 47–57.
- Espinal MR, Miller BW, Mora JM** (2021) First record of *Centronycteris centralis* (Chiroptera: Emballonuridae) from Honduras. *Notas sobre Mamíferos Sudamericanos* 3: e21.11.1. <http://doi.org/10.31687/saremNMS.21.11.1>
- Esquivel DA, Pereira MJR, Stuhler JD, Rossoni DM, Velazco PM, Bianchi FM** (2022) Multiple lines of evidence unveil cryptic diversity in the *Lophostoma brasiliense* (Chiroptera: Phyllostomidae) complex. *Systematics and Biodiversity* 20 (1): 1–21. <https://doi.org/10.1080/14772000.2022.2110172>
- Feijó A, Da Rocha PA, Althoff SL** (2015) New species of *Histiotus* (Chiroptera: Vespertilionidae) from northeastern Brazil. *Zootaxa* 4048 (3): 412–427. <https://doi.org/10.11646/zootaxa.4048.3.4>
- Gardner AL** (2008) Family Furipteridae Gray, 1866. In: Gardner AL (Ed.) *Mammals of South America, vol. 1, marsupials, xenarthrans, shrews, and bats*. University of Chicago Press, Chicago, USA, 389–391.
- Görföl T, Huang JC, Csorba G, Györössy D, Estók P, Kingston T, Szabadi KL, McArthur E, Senawi J, Furey NM, Tu VT, Thong VD, Khan FAA, Jinggong ER, Donnelly M, Kumaran JV, Liu J, Chen S, Tuanmu M, Ho Y, Chang H, Elias N, Abdullah N, Lim L, Squire CD, Zsebők S** (2022) ChiroVox: a public library of bat calls. *PeerJ* 10: e12445. <https://doi.org/10.7717/peerj.12445>
- Gregorin R, Moras LM, Acosta LH, Vasconcellos KL, Poma JL, dos Santos FR, Paca RC** (2016) A new species of *Eumops* (Chiroptera: Molossidae) from southeastern Brazil and Bolivia. *Mammalian Biology* 81 (3): 235–246. <https://doi.org/10.1016/j.mambio.2016.01.002>
- Handley CO Jr** (1976) *Mammals of the Smithsonian Venezuelan Project*. Brigham Young University Science Bulletin, Biological Series 20: 1–91.
- Harrison DL** (1975) *Macrophyllum macrophyllum*. *Mammalian Species* 62: 1–3. <https://doi.org/10.2307/3503986>
- Hintze F, Arias-Aguilar A, Dias-Silva L, Delgado-Jaramillo M, Rodrigues CS, Jucá T, Lyrio FM, Almeida M, Bezerra B, Aguiar LMS, Ramos MJP, Bernard E** (2020) Molossid unlimited: extraordinary extension of range and unusual vocalization patterns of the bat *Promops centralis*.

- Journal of Mammalogy 101 (2): 417–432. <https://doi.org/10.1093/jmammal/gyz167>
- Hood C, Gardner AL** (2008) Family Emballonuridae Gervais, 1856. In: Gardner AL (Ed.) Mammals of South America, vol. 1, marsupials, xenarthrans, shrews, and bats. University of Chicago Press, Chicago, USA, 188–207.
- Husson AM** (1962) The bats of Suriname. Zoölogische Verhandelingen (Rijksmuseum van Natuurlijke Historie, Leiden) 58: 1–282.
- Ibáñez C** (1985) Notas sobre distribución de quirópteros en Bolivia (Mammalia, Chiroptera). Historia Natural 5: 329–333.
- Ibáñez C, Ochoa J** (1989) New records of bats from Bolivia. Journal of Mammalogy 70 (1): 216–219. <https://doi.org/10.2307/1381694>
- Jones J, Hood C** (1993) Synopsis of South American bats of the Family Emballonuridae. Occasional Papers (The Museum, Texas Tech University) 155: 1–32.
- Jung K, Kalko EKV, Helversen VO** (2007) Echolocation calls in Central American emballonurid bats: signal design and call frequency alternation. Journal of Zoology 272: 125–137. <https://doi.org/10.1111/j.1469-7998.2006.00250.x>
- Jung K, Molinari J, Kalko EKV** (2014) Driving factors for the evolution of species-specific echolocation call design in new world free-tailed bats (Molossidae). PLoS ONE 9 (1): e85279. <https://doi.org/10.1371/journal.pone.0085279>
- Koopman KF** (1976) Zoogeography. In: Baker RJ, Jones JK, Carter DC (Eds.) Biology of bats of the New World family Phyllostomatidae. Part I. Special Publications (The Museum, Texas Tech University) 10: 39–47. <https://doi.org/10.5962/bhl.title.142603>
- LaVal RK** (1973) A revision of the Neotropical bats of the genus *Myotis*. Natural History Museum, Los Angeles County, Science Bulletin 15: 1–54.
- López-Vidal JC, Elizalde-Arellano C** (2006) Las colecciones mastozoológicas en la formación de recursos humanos en taxonomía y docencia. In: Lorenzo C, Espinoza E, Briones M, Cervantes FA (Eds.) Colecciones mastozoológicas de México. Instituto de Biología, Universidad Nacional Autónoma de México & Asociación Mexicana de Mastozología, A.C., Mexico City, Mexico, 123–134.
- Loureiro LO, Engstrom MD, Lim BK** (2020) Single nucleotide polymorphisms (SNPs) provide unprecedented resolution of species boundaries, phylogenetic relationships, and genetic diversity in the mastiff bats (*Molossus*). Molecular Phylogenetics and Evolution 143: 106690. <https://doi.org/10.1016/j.ympev.2019.106690>
- Ludeña JP, Medina CE** (2017) Segundo registro de *Cyttarops alecto* y *Eumops wilsoni* (Mammalia: Chiroptera) en el Perú. Revista Peruana de Biología 24 (4): 401–405. <https://doi.org/10.15381/rpb.v24i4.14065>
- Mantilla-Meluk H, Baker RJ** (2010) New species of *Anoura* (Chiroptera: Phyllostomidae) from Colombia, with systematic remarks and notes on the distribution of the *A. geoffroyi* complex. Occasional Papers (The Museum, Texas Tech University) 292: 1–19.
- Mantilla-Meluk H, Aguirre LF, Jiménez-Ortega AM** (2012) First record of *Anoura aequatoris* (Lönnberg, 1921) (Chiroptera: Phyllostomidae) from Bolivia. Revista Biodiversidad Neotropical 2 (1): 21–26.
- Mantilla-Meluk H, Siles L, Aguirre LF** (2014) Geographic and ecological amplitud in the nectarivorous bat *Anoura fistulata* (Phyllostomidae: Glossophaginae). Caldasia 36 (2): 373–388.
- Masson D, Cosson JF** (1992) *Cyttarops alecto* (Emballonuridae) et *Lasiurus castaneus* (Vespertilionidae), deux chiroptères nouveaux pour la Guyane Française. Mammalia 56 (3): 475–478.
- Morales-Martínez DM, López-Arévalo HF, Vargas-Ramírez M** (2021) Beginning the quest: phylogenetic hypothesis and identification of evolutionary lineages in bats of the genus *Micronycteris* (Chiroptera, Phyllostomidae). ZooKeys 1028: 135–159. <https://doi.org/10.3897/zookeys.1028.60955>
- Moratelli R, Dias D** (2015) A new species of nectar-feeding bat, genus *Lonchophylla*, from the Caatinga of Brazil (Chiroptera, Phyllostomidae). ZooKeys 514: 73–91. <https://doi.org/10.3897/zookeys.514.10013>
- Moratelli R, Wilson DE** (2014) A new species of *Myotis* (Chiroptera, Vespertilionidae) from Bolivia. Journal of Mammalogy 95 (4): E17–E25. <https://doi.org/10.1644/14-mamm-149>
- Moya IM, Galarza MI, Vargas A, Aguirre LF** (2007) Murciélagos de las Yungas de Bolivia. Biota, Cochabamba, Bolivia, 70 pp.
- Novaes R, Jacob G, Souza R, Sauwen C, Felix S, Avilla L** (2012) New record of *Furipterus horrens* (Cuvier, 1828) (Mammalia, Chiroptera) from the Cerrado of Tocantins state with a compilation of the known distribution within Brazil. Check List 8 (6): 1359–1361. <https://doi.org/10.15560/8.6.1359>
- Ochoa GJ, Soriano PJ, Hernandez-Camacho J** (1994) Sobre la presencia de *Cyttarops alecto* (Chiroptera: Emballonuridae) en Colombia. Trianea 5: 411–414.
- Ochoa GJ, O'Farrel MJ, Miller BW** (2000) The contribution of acoustic methods to the study of insectivorous bat diversity in protected areas from northern Venezuela. Acta Chiropterologica 2: 171–183.
- Ochoa GJ, Bevilacqua M, García FJ** (2005) Evaluación ecológica rápida de las comunidades de mamíferos en cinco localidades del Delta del Orinoco, Venezuela. Inter-ciencia 30: 466–475.
- Ochoa GJ, Sánchez J, García F, Clavijo JA** (2008) An undescribed behavior of the Isabelle's ghost bat (*Diclidurus isabellus*), with distributional notes in Venezuela. Memorias de la Fundación La Salle de Ciencias Naturales 170: 141–147.
- Ochoa GJ, García FJ, Caura S, Sánchez JH** (2009) Mamíferos de la Cuenca del Río Caura: listado taxonómico y distribución conocida. Memorias de la Fundación La Salle de Ciencias Naturales 170: 5–80.
- Ojeda RA, Barquez RM** (1978) Contribución al conocimiento de los quirópteros de Bolivia. Neotrópica 24: 33–38.
- Paca RC, Acosta LH, Aguanta FS** (2012) First record of *Tonatia bidens* (Spix, 1823) (Chiroptera: Phyllostomidae), in Bolivia. Chiroptera Neotropical 18 (1): 1063–1066.
- Pacheco V, Solari S, Velazco PM** (2004) A New species of *Carollia* (Chiroptera: Phyllostomidae) from the Andes of Peru and Bolivia. Occasional Papers, Museum of Texas

- Tech University 236: 1–15.
- Parlos JA, Timm RM, Swier VJ, Zeballos H, Baker RJ** (2014) Evaluation of the paraphyletic assemblages within Lonchophyllinae, with description of a new tribe and genus. *Occasional Papers, Museum of Texas Tech University* 320: 1–23.
- Pine RH** (1972) The bats of the genus *Carollia*. Technical Monograph, Agricultural Experimental Station, Texas A&M University 8: 1–125.
- Poma-Urey JL, Acosta LH, Ingala MR** (2019) Dos especies de *Eptesicus* Rafinesque, 1820 (Chiroptera, Vespertilionidae) registradas por primera vez en Bolivia. *Ecología en Bolivia* 54 (2): 155–161.
- Poma-Urey JL, Acosta L, Paca RC** (2020) Presencia de *Micronycteris sanborni* Simmons, 1996 (Chiroptera, Phyllostomidae) en Bolivia. *Kempffiana* 16 (1): 49–59.
- Poma-Urey JL, Acosta LH, Ingala MR, Revollo SG, Meza MA, Gutiérrez-Cruz S, Zabala-Pedraza JC, Peñaranda M, Salazar-Bravo J** (2021) Nueva especie de *Peropteryx* (Chiroptera: Emballonuridae) para Bolivia. *Notas sobre Mamíferos Sudamericanos* 3: e21.2.5. <https://doi.org/10.31687/saremnms.21.2.5>
- Porter CA, Hooper SR, Cline CA, Hoffmann FG, Baker RJ** (2007) Molecular phylogenetics of the phyllostomid bat genus *Micronycteris* with descriptions of two new subgenera. *Journal of Mammalogy* 88: 1205–1215. <https://doi.org/10.1644/06-mamm-a-292r.1>
- Ruedas LA, Salazar-Bravo J, Dragoo JW, Yates TL** (2000) The importance of being earnest: what, if anything, constitutes a “specimens examined?” *Molecular Phylogenetics and Evolution* 17: 129–132. <https://doi.org/10.1006/mpev.2000.0737>
- Salazar-Bravo J, Tarifa T, Aguirre LF, Yensen E, Yates TL** (2003) Revised checklist of Bolivian Mammals. *Occasional Papers, Museum of Texas Tech University* 220: 1–27.
- Sampaio E, Lim B, Peters S** (2016) *Centronycteris maxmilianii*. The IUCN Red List of Threatened Species 2016: e.T4112A22002444. <https://doi.org/10.2305/iucn.uk.2016-2.rlts.t4112a22002444.en>. Accessed on: 2022–11–30.
- Sanborn CC** (1932a) Neotropical bats in the Carnegie Museum. *Annals of the Carnegie Museum* 21: 171–183.
- Sanborn CC** (1932b) The bats of the genus *Eumops*. *Journal of Mammalogy* 13 (4): 347–357. <https://doi.org/10.2307/1374140>
- Sanborn CC** (1933) Bats of the genera *Anoura* and *Lonchoglossa*. *Field Museum of Natural History, Zoological Series* 20: 23–28.
- Sanborn CC** (1937) American bats of the subfamily Emballonurinae. *Field Museum of Natural History, Zoological Series* 20: 321–354.
- Sánchez-Hernández C, Romero-Almaraz ML, Schnell GD** (2005) New species of *Sturnira* (Chiroptera: Phyllostomidae) from northern south America. *Journal of Mammalogy* 86 (5): 866–872. [https://doi.org/10.1644/1545-1542\(2005\)86\[866:nsoscp\]2.0.co;2](https://doi.org/10.1644/1545-1542(2005)86[866:nsoscp]2.0.co;2)
- Schnitzler HU, Kalko EKV** (2001) Echolocation by insect-eating bats. *BioScience* 51: 557–569.
- Semedo TBF, Feijó A** (2016) Filling the gap: first record of the transparent-winged big-eared bat *Histiotus diaphanopterus* (Chiroptera: Vespertilionidae) in southwestern Brazil. *Mammalia* 81 (3): 323–327. <https://doi.org/10.1515/mammalia-2016-0007>
- Siles L** (2007) Noteworthy records of *Eptesicus chiriquinus* and *Eptesicus andinus* (Vespertilionidae) from Bolivia. *Bat Research News* 48 (2): 31–33.
- Siles L, Brooks DM, Aranibar H, Tarifa T, Vargas JM, Rojas JM, Baker RJ** (2013) A new species of *Micronycteris* (Chiroptera: Phyllostomidae) from Bolivia. *Journal of Mammalogy* 94: 881–896. <https://doi.org/10.1644/12-mamm-a-259.1>
- Siles L, Hidalgo-Cossio M, Amusquivar-Calustro A, Camacho T, Huayta-Sarzuri G** (2021) First record of *Peropteryx pallidoptera* (Chiroptera: Emballonuridae) in Bolivia. *Therya Notes* 2: 51–55. https://doi.org/10.12933/therya_notes-21-35
- Siles L, Wallace RB** (2021) First record of the rare bat *Gardnerycteris koepckeae* (Gardner & Patton, 1972) (Chiroptera, Phyllostomidae) in Bolivia. *Check List* 17 (1): 1–6. <https://doi.org/10.15560/17.1.1>
- Simmons NB, Cinarrello AL** (2023) Bat species of the world: a taxonomic and geographic database. Version 1.3. <https://batnames.org/query.html>. Accessed on: 2023-5-24.
- Simmons NB, Voss RS** (1998) The mammals of Paracou, French Guiana, a Neotropical lowland rainforest fauna. Part 1. *Bulletin of the American Museum of Natural History* 237: 1–219.
- Solari S, Baker RJ** (2006) Mitochondrial DNA sequence, karyotypic, and morphological variation in the *Carollia castanea* species complex (Chiroptera: Phyllostomidae) with description of a new species. *Occasional Papers, Museum of Texas Tech University* 254: 1–16.
- Solari S** (2015) *Saccopteryx canescens*. The IUCN Red List of Threatened Species 2015: e.T19805A22005456. <https://doi.org/10.2305/iucn.uk.2015-4.rlts.t19805a22005456.en>. Accessed on: 2022-11-4.
- Solari S** (2019) *Molossus pretiosus*. The IUCN Red List of Threatened Species 2019: e.T13649A22106312. <https://doi.org/10.2305/iucn.uk.2019-1.rlts.t13649a22106312.en>. Accessed on: 2022–11–10.
- Solari S, Medellín R, Rodríguez-Herrera B, Tavares VC, Garbino GST, Camacho MA, Tirira D, Lim B, Arroyo-Cabrales J, Rodríguez-Durán A, Dumont E, Burneo S, Aguirre LF, Tschapka M, Espinosa D** (2019) Family Phyllostomidae (New World leaf-nosed bats). In: Wilson DE, Mittermeier RA (Eds.) *Handbook of the mammals of the world, bats, volume 9*. Lynx Edicions, Barcelona, Spain, 444–583.
- Starrett A, de la Torre L** (1964) Notes on a collection of bats from Central America, with the third record for *Cyttarops alecto* Thomas. *Zoologica* 49 (1): 53–63.
- Starrett A, Casebeer RS** (1968) Records of bats from Costa Rica. *Los Angeles Natural History Museum Contribution to Science* 148: 1–21.
- Taddei VA, Uieda W** (2001) Distribution and morphometrics of *Natalus stramineus* from South America (Chiroptera, Natalidae). *Iheringia, Serie Zoologia* 91: 123–132. <https://doi.org/10.1590/S0073-47212001000200018>
- Tarifa T** (2014) Historia de la Mastozoología en Bolivia. In: Ortega J, Martínez JL, Tirira D (Eds.) *Historia de la mastozoología en Latinoamérica, las Guayanas y el Caribe*.

- Editorial Murciélago Blanco y Asociación Ecuatoriana de Mastozoología, Quito y México, DF., 69–106.
- Tavares VD, Estefano P, Bobrowiec D, Farias SG** (2012) First record of the rare bat *Cyttarops alecto* (Thomas, 1913) (Chiroptera: Emballonuridae) for the western Brazilian Amazonia, with comments on the type locality. *Mammalia* 76: 345–349. <https://doi.org/10.1515/mammalia-2011-0117>
- Taylor PJ, Lim BK, Pennay M, Soisook P, Loureiro LO, Moras LM, Kingston T** (2019). Familia Molossidae. In: Wilson DE, Mittermeier RA (Eds.) Handbook of the mammals of the world, bats, volume 9. Lynx Edicions. Barcelona, Spain, 598–672.
- Tejedor A** (2003) First record of *Saccopteryx canescens* (Chiroptera: Emballonuridae) for southeastern Peru. *Chiroptera Neotropical* 9: 162–164.
- Terán M** (2010) Nuevos registros de murciélagos (Chiroptera: Phyllostomidae: Phyllostominae) para el departamento de La Paz, Bolivia. *Revista Boliviana de Ecología y Conservación Ambiental* 27: 85–91.
- Thomas O** (1902) On mammals from Cochabamba, Bolivia, and the region north of that place. *Annals and Magazine of Natural History (series 7)* 9: 125–143.
- Thomas O** (1913) On some rare Amazonian mammals from the collection of the Para Museum. *Annals and Magazine of Natural History (series 8)* 12: 130–136.
- Thomas O** (1916) Notes on the bats of the genus *Histiotus*. *Annals and Magazine of Natural History (series 8)* 17: 272–276.
- Uieda W, Sazima I, Storti-Filho A** (1980) Aspectos da biologia do morcego *Furipterus horrens*. *Revista Brasileira de Biologia* 40 (1): 59–66.
- Vargas-Espinoza A** (2007) Familia Emballonuridae (Gervais, 1855). In: Aguirre LF (Ed.) Historia natural, distribución y conservación de los murciélagos de Bolivia. Fundación Simón I. Patiño, Santa Cruz, Bolivia, 158–173.
- Vargas A, Balderrama JA** (2009) Primer registro de *Artibeus (Koopmania) concolor* (Chiroptera: Stenodermatinae) para Bolivia. *Mastozoología Neotropical* 16 (2): 495–497.
- Velazco PM** (2005) Morphological phylogeny of the bat genus *Platyrrhinus* (Chiroptera: Phyllostomidae) with the description of four new species. *Fieldiana Zoology* 105: 1–53. [https://doi.org/10.3158/0015-0754\(2005\)105\[1:mpo](https://doi.org/10.3158/0015-0754(2005)105[1:mpo)
- [tbg\]2.0.co;2](https://doi.org/10.3158/0015-0754(2005)105[1:mpo)
- Velazco S, Pacheco V, Meschede A** (2011) First occurrence of the rare emballonurid bat *Cyttarops alecto* (Thomas, 1913) in Peru—only hard to find or truly rare? *Mammalian Biology* 76: 373–376. <https://doi.org/10.1016/j.mambio.2010.03.006>
- Velazco PM, Guevara L, Molinari J** (2018) Systematics of the broad-nosed bats, *Platyrrhinus umbratus* (Lyon, 1902) and *P. nigellus* (Gardner and Carter, 1972) (Chiroptera: Phyllostomidae), based on genetic, morphometric, and ecological niche analyses. *Neotropical Biodiversity* 4 (1): 119–133. <https://doi.org/10.1080/23766808.2018.1494481>
- Velazco PM, Patterson BD** (2019) Small mammals of the Mayo River Basin in northern Peru, with the description of a new species of *Sturnira* (Chiroptera: Phyllostomidae). *Bulletin of the American Museum of Natural History* 429: 1–67. <https://doi.org/10.1206/0003-0090.429.1.1>
- Velazco PM, Voss RS, Fleck DW, Simmons NB** (2021) Mammalian diversity and Matses ethnomammalogy in Amazonian Peru, part 4: bats. *Bulletin of the American Museum of Natural History* 451 (1): 1–200. <https://doi.org/10.1206/0003-0090.451.1.1>
- Venzal JM, Nava S, Terassini FA, Ogrzewalska M, Camargo LMA, Labruna MB** (2013) *Ornithodoros peropteryx* (Acari: Argasidae) in Bolivia: an argasid tick with a single nymphal stage. *Experimental and Applied Acarology* 61: 231–241. <https://doi.org/10.1007/s10493-013-9689-3>
- Voss RS, Emmons LH** (1996) Mammalian diversity in Neotropical lowland rainforests: a preliminary assessment. *Bulletin of the American Museum of Natural History* 230: 1–115.
- Webster WD, Jones JK** (1980) Noteworthy records of bats from Bolivia. *Occasional Papers, Museum of Texas Tech University* 68: 1–7 6.
- Wilson DE, Salazar JA** (1990) Los murciélagos de la Reserva de la Biosfera “Estación Biológica Beni”. *Ecología en Bolivia* 13: 47–56.
- Zortea M, Sampaio E, Lim B, Peters S** (2016) *Glyphonycteris behnii*. The IUCN Red List of Threatened Species 2016: e.T13375A22130995. <https://doi.org/10.2305/iucn.uk.2016-2.rlts.t13375a22130995.en>. Accessed on: 2022–11–10.